

Domain Structure of Black Hole Space-Times

Troels Harmark

The Niels Bohr Institute

Blegdamsvej 17, 2100 Copenhagen Ø, Denmark

harmark@nbi.dk

Abstract

We introduce the domain structure for stationary black hole space-times. Given a set of commuting Killing vector fields of the space-time the domain structure lives on the submanifold where at least one of the Killing vector fields have zero norm. Depending on which Killing vector field has zero norm the submanifold is naturally divided into domains. A domain corresponds either to a set of fixed points of a spatial symmetry or to a Killing horizon, depending on whether the characterizing Killing vector field is space-like or time-like near the domain. The domain structure provides invariants of the space-time, both topological and geometrical. It is defined for any space-time dimension and any number of commuting Killing vector fields. We examine the domain structure for asymptotically flat space-times and find a canonical form for the metric of such space-times. The domain structure generalizes the rod structure introduced for space-times with $D - 2$ commuting Killing vector fields. We analyze in detail the domain structure for Minkowski space, the Schwarzschild-Tangherlini black hole and the Myers-Perry black hole in six and seven dimensions. Finally we consider the possible domain structures for asymptotically flat black holes in six and seven dimensions.

Contents

1	Introduction and summary	1
2	Canonical form of metric	3
2.1	Preliminaries	3
2.2	Canonical form for particular class of metrics	5
2.3	Canonical form for general class of metrics	6
3	Domain structure	7
3.1	The flow of the Killing vector fields	7
3.2	Submanifolds and the kernel of G	8
3.3	Domains and their directions	10
3.4	Reduction to the rod structure for $n = 2$	12
4	Domain structure of six dimensional black holes	13
5	Domain structure of seven dimensional black holes	16
6	Possible new domain structures in six and seven dimensions	20
6.1	Six-dimensional asymptotically flat space-times	20
6.2	Seven-dimensional asymptotically flat space-times	23
7	Discussion and outlook	26
A	Einstein equations for space-times with Killing vector fields	27
B	Parameterizations of topologies from domain structure	29
	References	31

1 Introduction and summary

It has been realized in recent years that the dynamics of black holes in dimension $D \geq 5$ is much richer than in four dimensions. In four dimensions the famous uniqueness theorems [1] state that given the asymptotic charges, *i.e.* the mass, angular momentum and the electric and magnetic charges, there is at most one available black hole solution, namely the Kerr-Newman solution. In dimensions $D \geq 5$ there are instead a number of available solutions given the asymptotic charges, as first realized with the discovery of the black ring in five dimensions [2]. This naturally brings up the question of whether one can find a general set of invariants, in addition to the asymptotic charges, that characterize a black hole space-time for $D \geq 5$.

In this paper we propose a set of invariants given a stationary black hole space-time with any number of space-time dimensions and any number of commuting Killing vector fields. We call this set of invariants the *domain structure*. The domain structure lives on the submanifold where at least one of the Killing vector fields have zero norm. Depending on which Killing vector field has zero norm the submanifold is naturally divided into domains. A domain corresponds either to a set of fixed points of a spatial symmetry or to a Killing horizon,

depending on whether the characterizing Killing vector field is space-like or time-like near the domain.

The domain structure generalizes the so-called *rod structure* proposed in [3] as the set of invariants characterizing asymptotically flat black holes in five dimensions which are solutions of vacuum Einstein equations and possess two rotational Killing vector fields. For this class of solutions the submanifold for which at least one of the Killing vector fields have zero norm can in certain canonical coordinates be seen as a line. This line is then divided into intervals called rods according to which Killing vector field has a zero norm. Such rods of five-dimensional black holes was first considered for generalized Weyl solutions in [4]. The proposal of [3] that the rod structure provides a characterization of asymptotically flat black holes in five dimensions which are solutions of vacuum Einstein equations and possess two rotational Killing vector fields is supported by the uniqueness theorem of [5] which states that a black hole space-time with a single event horizon is unique given the rod structure and the asymptotic charges.

The domain structure provides in particular a generalization of the rod structure for the case of five-dimensional black hole space-times with two rotational Killing vector fields (and more generally solutions with $D - 2$ commuting linearly independent Killing vector fields) since in the approach of this paper we can analyze solutions with matter fields such as gauge fields and scalar fields. This overlaps with previous generalizations of the rod structure [6]. We reproduce furthermore the constraints on the rod structure derived in [5].

The domain structure provides invariants of the black hole space-time, both topological and geometrical. It reveals certain aspects of the global structure of the black hole space-time. In particular one can read off the topology of the event horizon(s). It can also help in exploring what black hole space-times are possible. The more commuting Killing vector fields one has, the more invariants one obtains and the more constraints one finds on the possible black hole space-times. In terms of topology of the event horizon the topological censorship theorem says that it should be of positive Yamabe type (i.e. it must admit a metric of positive curvature) [7]. If we have more than just the Killing vector field associated with stationarity of the space-time we can give further restrictions on the topology. Thus, not only we can provide invariants to characterize the space-time, we can also use the domain structure to provide limitations on what types of black holes are possible. In addition to the topological invariants the geometrical invariants which measures the volumes of each domain can be used as a further characterization of the space-time.

We will always assume that our black hole space-time is stationary, *i.e.* it has a Killing vector field which is time-like far away from the event horizon(s). If this is the only Killing vector field the domain structure invariants will coincide with the previously known topological data and physical parameters of the black hole space-time. For example, the domain structure will give the topology and the area of the event horizon(s). Given any number of additional (asymptotically spatial) commuting Killing vector fields one finds new invariants of the black hole space-time. For asymptotically flat space-times the existence of at least one rotational Killing vector field is guaranteed by the rigidity theorems of [8].

We assume in this paper that the black hole space-times are either asymptotically flat or

asymptotically Kaluza-Klein space (here defined as a $(D - q)$ -dimensional Minkowski space times a q -dimensional torus). Under this assumption we can find a canonical form of the metric which is used to define the domain structure. However, the general analysis of this paper can also work for space-times with other asymptotics, such as asymptotically Anti-de Sitter space-times. We save this generalization for a future publication [9].

More concretely the paper is built up as follows. In Section 2 we find a canonical form of the metric for all asymptotically flat black hole space-times and all space-times which are asymptotically Kaluza-Klein space.

In Section 3 we define the domain structure for black hole space-times. We analyze the structure of the kernel of the metric on the commuting Killing vector fields and how this gives rise to a hierarchy of submanifolds. We find that the submanifold corresponding to the zero norm Killing vector fields is naturally divided into domains and use this to define the domain structure. We end with considering the special case of $D - 2$ commuting Killing vector fields where we obtain the rod structure now for a more general class of solutions.

In Sections 4 and 5 we analyze the Minkowski space, the Schwarzschild-Tangherlini black hole and the Myers-Perry black hole [10] in six and seven dimensions. These space-times all possess $D - 3$ commuting Killing vector fields. We find coordinates such that the metric is put in a canonical form. Using these coordinates we find the domain structure of the space-times.

Finally we consider in Section 6 which domain structures are possible for asymptotically flat black hole space-times in six and seven dimensions with $D - 3$ commuting Killing vector fields. Here we analyze in particular the domain structures for the new types of black holes found recently using the Blackfold approach [11, 12]. We also consider the static numerical solutions recently found in [13].

We end in Section 7 with discussing the implications of the results of this paper and what new directions we can take. In particular we discuss whether the domain structure along with the asymptotic charges give enough invariants to fully characterize an asymptotically flat black hole space-time. In line with this we conjecture a uniqueness theorem for a certain class of black hole space-times.

2 Canonical form of metric

We consider in the following a given D -dimensional space-time with p commuting linearly independent Killing vector fields. In detail we are given a D -dimensional manifold $\mathcal{M}_D$ with a Lorentzian signature metric with p commuting linearly independent Killing vector fields $V_{(i)}$, $i = 0, 1, \dots, p - 1$. The Killing vector fields are such that they generate the isometry group $\mathbb{R} \times U(1)^{p-1}$. In particular the $p - 1$ $U(1)$ symmetries are generated by the $p - 1$ space-like Killing vector fields $V_{(i)}$, $i = 1, \dots, p - 1$, while the Killing vector field $V_{(0)}$ generates the $\mathbb{R}$ isometry. In the following we present the canonical form of the metric for such a space-time.

2.1 Preliminaries

As stated above we are given a D -dimensional space-time $\mathcal{M}_D$ with p commuting linearly independent Killing vector fields $V_{(i)}$, $i = 0, 1, \dots, p - 1$. Define $n = D - p$. We can always find

a coordinate system $x^0, x^1, \dots, x^{p-1}, y^1, \dots, y^n$ such that

$$V_{(i)} = \frac{\partial}{\partial x^i} \quad (2.1)$$

for $i = 0, 1, \dots, p-1$. We then write the metric as

$$ds^2 = G_{ij}(dx^i + A_a^i dy^a)(dx^j + A_b^j dy^b) + \tilde{g}_{ab} dy^a dy^b \quad (2.2)$$

with $i, j = 0, 1, \dots, p-1$ and $a, b = 1, \dots, n$, and where G_{ij} , A_a^i and $\tilde{g}_{ab}$ only depend on y^a . In Appendix A we compute the Ricci tensor for the metric (2.2).

The goal of this paper is to understand the structure of the kernel of the metric G_{ij} on the commuting Killing vector fields for black hole space-times. If we have a point q for which the kernel $\ker G(q)$ is non-trivial it means that there exists (at least one) Killing vector field W , which is a (non-zero) linear combination of $V_{(i)}$, $i = 0, 1, \dots, p-1$, such that $G(q)W = 0$. Equivalently, one can say that the norm of W , as measured with the metric G_{ij} , is zero at q . Thus, the submanifold where at least one linear combination of the Killing vector fields $V_{(i)}$ has zero norm corresponds to the points where the kernel $\ker G$ is non-trivial.

It is clear from this that whenever we have a point q for which $\ker G(q)$ is non-trivial we have that $\det G(q) = 0$. We therefore use the function $\det G$ on the space-time to define one of the coordinates in our canonical form for the metric.

Define now the n -dimensional manifold $\mathcal{N}_n$ as the quotient space $\mathcal{M}_D / \sim$ where the equivalence relation $\sim$ is such that two points in $\mathcal{M}_D$ are equivalent if they can be connected by an integral curve of a linear combination of the Killing vector fields $V_{(i)}$, $i = 0, 1, \dots, p-1$. This is known as the *orbit space* of the space-time since each point in $\mathcal{N}_n$ corresponds to an orbit of the $\mathbb{R} \times U(1)^{p-1}$ symmetry of the commuting Killing vector fields. The n -dimensional manifold $\mathcal{N}_n$ is naturally equipped with the n -dimensional part of the metric (2.2)

$$ds_n^2 = \tilde{g}_{ab} dy^a dy^b \quad (2.3)$$

where we sum over $a, b = 1, 2, \dots, n$.

On the n -dimensional manifold $\mathcal{N}_n$ the fields A_a^i , $i = 0, 1, \dots, p-1$, can be thought of as components of p $U(1)$ gauge fields. Consider the coordinate transformation $x^i \rightarrow x^i - \alpha^i(y^a)$ with y^a kept fixed. Under this coordinate transformation the Killing vectors are still of the form (2.1) and for the metric (2.2) G_{ij} and $\tilde{g}_{ab}$ stays the same while A_a^i transforms as $A_a^i \rightarrow A_a^i + \partial \alpha^i / \partial y^a$. We see that this is a gauge transformation of the $U(1)$ gauge field $A^i = A_a^i dy^a$.

In the following we would like to define new coordinates on the n -dimensional manifold $\mathcal{N}_n$ with metric (2.3) suitable for examining the kernel of the metric G_{ij} on the commuting Killing vector fields $V_{(i)}$. To this end, define the function $r(y^a)$ on $\mathcal{N}_n$ as

$$r^m \equiv \sqrt{|\det G_{ij}|} \quad (2.4)$$

for a positive real number m . We see then that the kernel of G_{ij} corresponds to $r = 0$. We assume that $(\partial r / \partial y^1, \dots, \partial r / \partial y^n) \neq 0$ on $\mathcal{N}_n$ up to a subspace of n -volume zero. Define now the vector field $\chi = \chi^a \partial / \partial y^a$ on $\mathcal{N}_n$ by $\chi^a = \tilde{g}^{ab} (\partial r / \partial y^b)$. Define the equivalence relation $\sim$

on $\mathcal{N}_n$ such that two points are equivalent if they are connected by an integral curve of χ . We can then consider the quotient space $\mathcal{N}_n/\sim$, which is an $(n-1)$ -dimensional manifold. Consider coordinates $z^1, \dots, z^{n-1}$ on $\mathcal{N}_n/\sim$. We can extend these coordinates to functions on $\mathcal{N}_n$. We see then that z^α is constant on the integral curves of χ ($\alpha = 1, 2, \dots, n-1$). Therefore, $\chi^\alpha(\partial z^\alpha/\partial y^a) = 0$. Clearly $r, z^1, \dots, z^{n-1}$ is a coordinate system on $\mathcal{N}_n$ and $g^{rz^\alpha} = 0$, $\alpha = 1, 2, \dots, n-1$. We can thus write the metric on $\mathcal{N}_n$ as

$$ds_n^2 = e^{2A} dr^2 + \sum_{\alpha, \beta=1}^{n-1} \tilde{\Lambda}_{\alpha\beta} dz^\alpha dz^\beta \quad (2.5)$$

where $A(r, z^\alpha)$ and $\tilde{\Lambda}_{\alpha\beta}(r, z^\gamma)$ are functions.

2.2 Canonical form for particular class of metrics

Before treating the general class of metrics we first consider a special class of metrics. This class consists of metrics solving the vacuum Einstein equations $R_{\mu\nu} = 0$. For a metric of the form (2.2) these equations can be written as (A.4)-(A.6) in Appendix A. We demand in addition that the p Killing vector fields are such that

$$V_{(0)}^{[\mu_1} V_{(1)}^{\mu_2} \dots V_{(p-1)}^{\mu_p} D^\nu V_{(i)}^{\rho]} = 0 \quad \text{for all } i = 0, 1, \dots, p-1 \quad (2.6)$$

Among the space-times in this class is D -dimensional Minkowski space, which in addition to the time-translation Killing vector field has $[(D-1)/2]$ rotational Killing vector fields so that we can take $p \leq 1 + [(D-1)/2]$.¹ Also the Kaluza-Klein space $\mathbb{R}^{1, D-1-q} \times T^q$ is in this class with $p \leq 1 + q + [(D-q-1)/2]$.

By Theorem A.1 of Appendix A the condition (2.6) means we can write the metric as

$$ds^2 = G_{ij} dx^i dx^j + \tilde{g}_{ab} dy^a dy^b \quad (2.7)$$

where we sum over $i, j = 0, 1, \dots, p-1$ and $a, b = 1, 2, \dots, n$ (*i.e.* we have $D = p + n$) and where $V_{(i)} = \partial/\partial x^i$, $i = 0, 1, \dots, p-1$. We see that this corresponds to the metric (2.2) with $A_a^i = 0$. If we take the trace of Eq. (A.4) (with $A_a^i = 0$) we get $\partial_a(\sqrt{\tilde{g}} \tilde{g}^{ab} \partial_b r^m) = 0$ where $r(y^a)$ is defined in (2.4). In the (r, z^α) coordinate system introduced in Section 2.1 this gives $-\partial_r A + \partial_r \log \sqrt{\det \tilde{\Lambda}_{\alpha\beta}} + (m-1)/r = 0$. We see that it is natural to choose $m = 1$. Furthermore, we define $\nu \equiv (n-1)A$ and $\Lambda_{\alpha\beta} \equiv \exp(\frac{2A}{n-1}) \tilde{\Lambda}_{\alpha\beta}$. With this the trace equation $\partial_a(\sqrt{\tilde{g}} \tilde{g}^{ab} \partial_b r^m) = 0$ becomes

$$\partial_r \lambda = 0, \quad \lambda \equiv \sqrt{|\det \Lambda_{\alpha\beta}|} \quad (2.8)$$

One can show that D -dimensional Minkowski space $\mathbb{R}^{1, D-1}$ and also the Kaluza-Klein space $\mathbb{R}^{1, D-1-q} \times T^q$ admit z^α coordinates such that $\lambda = 1$. We show this explicitly for six and seven dimensional Minkowski space in Sections 4 and 5. Since for this particular class of metrics we are only interested in asymptotically flat solutions, or solutions that asymptote to

¹Of course D -dimensional Minkowski space has D commuting Killing vector fields but since our purpose here is to study black hole space-times we are only interested in the rotational and the time-translation Killing vector fields since these are the only ones that can be shared by asymptotically flat black hole solutions.

Kaluza-Klein space, we can demand without loss of generality that $\lambda \rightarrow 1$ for $r \rightarrow \infty$ with z^α fixed. Hence from (2.8) it follows that $\lambda = 1$ everywhere.

In conclusion, we get that for any D -dimensional space-time solving the vacuum Einstein equations with p commuting linearly independent Killing vector fields obeying (2.6) the metric can be put in the form

$$ds^2 = G_{ij}dx^i dx^j + e^{2(n-1)\nu} dr^2 + e^{2\nu} \Lambda_{\alpha\beta} dz^\alpha dz^\beta, \quad r^2 = |\det G_{ij}|, \quad \lambda = 1 \quad (2.9)$$

with the Killing vector fields given by (2.1). We call (2.9) the *canonical form of the metric* for this class of space-times.

2.3 Canonical form for general class of metrics

We now treat the general case. Thus, we consider here D -dimensional space-times with p commuting linearly independent Killing vector fields $V_{(i)}$, $i = 0, 1, \dots, p-1$. The Killing vector fields are such that they generate the isometry group $\mathbb{R} \times U(1)^{p-1}$.

Asymptotic flatness or Kaluza-Klein space asymptotics

From Section 2.1 we have that we can write the metric as in Eq. (2.2). The $\mathcal{N}_n$ part of the metric can furthermore be written as in Eq. (2.5) with r defined in (2.4). Choosing $m = 1$ and defining $\nu \equiv (n-1)A$ and $\Lambda_{\alpha\beta} \equiv \exp(\frac{2A}{n-1})\tilde{\Lambda}_{\alpha\beta}$ we see that the metric on $\mathcal{N}_n$ takes the form $e^{2(n-1)\nu} dr^2 + e^{2\nu} \Lambda_{\alpha\beta} dz^\alpha dz^\beta$ with $r^2 = |\det G_{ij}|$. Assuming furthermore that the space-time which we are considering asymptote to either D -dimensional Minkowski space or Kaluza-Klein space $\mathbb{R}^{1,D-1-q} \times T^q$ we can demand that $\lambda \rightarrow 1$ for $r \rightarrow \infty$. Thus, we can write the metric in the form

$$ds^2 = G_{ij}(dx^i + A^i)(dx^j + A^j) + e^{2(n-1)\nu} dr^2 + e^{2\nu} \Lambda_{\alpha\beta} dz^\alpha dz^\beta \quad (2.10)$$

$$r^2 = |\det G_{ij}|, \quad \lambda \rightarrow 1 \text{ for } r \rightarrow \infty$$

with the Killing vector fields given by (2.1) and $\lambda \equiv \sqrt{|\det \Lambda_{\alpha\beta}|}$. We call (2.10) the *canonical form of the metric* for solutions asymptoting to D -dimensional Minkowski space or Kaluza-Klein space $\mathbb{R}^{1,D-1-q} \times T^q$. This is a generalization of the form (2.9) for solutions of the vacuum Einstein equations obeying the condition (2.6). Instead here we can consider solutions which couples to any type of matter fields, such as gauge fields or scalar fields, since we do not use Einstein equations in getting the metric (2.10).

If we consider transforming from the coordinates y^a in (2.2) to two different coordinate systems (r, z^α) and $(\tilde{r}, \tilde{z}^\alpha)$ both making the metric to be of the canonical form (2.10) we immediately see from $r^2 = |\det G_{ij}|$ that $r(y^a) = \tilde{r}(y^a)$ and hence $\nu(y^a) = \tilde{\nu}(y^a)$. Considering now the transformation from (r, z^α) to $(\tilde{r}, \tilde{z}^\alpha)$ we see that $\tilde{r}(r, z^\alpha) = r$. In general we can write $\tilde{z}^\alpha(r, z^\beta)$. However, since $g^{rz^\alpha} = 0$ we have that $g^{\tilde{r}\tilde{z}^\alpha} = e^{-2(n-1)\nu}(\partial \tilde{z}^\alpha / \partial r)$. Therefore $\tilde{z}^\alpha$ cannot depend on r so the most general transformation is $\tilde{z}^\alpha(z^\beta)$. Imposing furthermore that $\lambda \rightarrow 1$ for $r \rightarrow \infty$ we get that *the only left over coordinate transformations in the canonical form (2.10) are $(n-1)$ -volume preserving diffeomorphisms of the z^α coordinates* (apart from rigid rotations of the x^i coordinates and the gauge transformations of A^i).

Other types of asymptotics

We can generalize the above to include metrics which are not asymptotically Minkowski space or Kaluza-Klein space. In general we imagine having a background space-time $\mathcal{M}_D^{(0)}$ that a given class of space-times asymptotes to. For this background space-time we can now find a function $\lambda_0(r, z^\alpha)$ such that $\lambda/\lambda_0 \rightarrow 1$ for $r \rightarrow \infty$. So for any space-time which asymptotes to $\mathcal{M}_D^{(0)}$ we can write the metric on the form

$$ds^2 = G_{ij}(dx^i + A^i)(dx^j + A^j) + e^{2(n-1)\nu} dr^2 + e^{2\nu} \Lambda_{\alpha\beta} dz^\alpha dz^\beta$$

$$r^2 = |\det G_{ij}|, \quad \frac{\lambda}{\lambda_0} \rightarrow 1 \text{ for } r \rightarrow \infty \quad (2.11)$$

In this way we can for example treat asymptotically Anti-de Sitter space-times. This will be considered in detail elsewhere [9]. For asymptotically de Sitter space-times the analysis proceeds differently since the asymptotic region includes the cosmological horizon for which $r = 0$ [9].

3 Domain structure

In this section we introduce the domain structure for black hole space-times. We focus here on asymptotically flat space-times and asymptotically Kaluza-Klein space-times. This means the metric can be put in the canonical form (2.10). The analysis is straightforwardly generalizable to other classes of space-times as well.

We consider in the following a D -dimensional manifold $\mathcal{M}_D$ with a Lorentzian signature metric with p commuting linearly independent Killing vector fields $V_{(i)}$, $i = 0, 1, \dots, p-1$. The Killing vector fields are such that they generate the isometry group $\mathbb{R} \times U(1)^{p-1}$. In particular the $p-1$ $U(1)$ symmetries are generated by the $p-1$ space-like Killing vector fields $V_{(i)}$, $i = 1, \dots, p-1$, while the Killing vector field $V_{(0)}$ generates the $\mathbb{R}$ isometry.

For purposes of our analysis we assume below that the following two regularity conditions on the metric (2.10) are obeyed: The A_a^i fields and the scalar fields $V_{(i)}^\mu V_{(j)}^\nu R_{\mu\nu}$ do not go to infinity for $r \rightarrow 0$.

This Section is built up as follows. In Section 3.1 we consider the flow of the p Killing vector fields. In Section 3.2 we examine how the kernel of the Killing metric gives rise to a natural hierarchy of submanifolds of $\mathcal{N}_n$. In Section 3.3 we define the domains and their directions and use this to define the domain structure. Finally in Section 3.4 we discuss the reduction of the domain structure to the rod structure in the special case of $n = 2$.

The analysis of this section builds on generalizations of methods used in Refs. [3] and [5].

3.1 The flow of the Killing vector fields

Before considering the form of the metric we consider here the flow of the Killing vector fields. This will be of importance for the analysis below. We define the flow of $V_{(i)}$ as

$$\sigma_s^{(i)}(x^0, \dots, x^i, \dots, x^{p-1}, y^1, \dots, y^n) = (x^0, \dots, x^i + s, \dots, x^{p-1}, y^1, \dots, y^n) \quad (3.1)$$

for $i = 0, 1, \dots, p-1$. For $i = 1, 2, \dots, p-1$ the Killing vector field $V_{(i)}$ generates a $U(1)$ isometry hence the flow is periodic. We normalize the periods of the flows with $i = 1, 2, \dots, p-1$ to be 2π . Note that for $i = 1, 2, \dots, p-1$ the Killing vector fields $V_{(i)}$ can not be time-like anywhere since then one would have closed time-like curves.

The set of Killing vector fields $V_{(i)}$, $i = 0, 1, \dots, p-1$, corresponds to a particular choice of basis. We are not entirely free to choose any basis. Consider the space-like Killing vectors $V_{(i)}$, $i = 1, \dots, p-1$. A new basis $W_{(i)}$, $i = 1, \dots, p-1$ is in general a linear combination

$$W_{(i)} = \sum_{j=1}^{p-1} U_{ij} V_{(j)} \quad (3.2)$$

Considering in particular $W_{(1)}$ this generates the flow

$$\sigma_s(x^0, x^1, x^2, \dots, x^{p-1}, y^1, \dots, y^n) = (x^0, x^1 + U_{11}s, x^2 + U_{12}s, \dots, x^{p-1} + U_{1,p-1}s, y^1, \dots, y^n) \quad (3.3)$$

We want $W_{(1)}$ to generate a $U(1)$ isometry and we choose the period of the flow to be 2π . From (3.3) we see this means that the U_{1i} entries should be relatively prime numbers. Thus we get the general requirement that $U \in GL(p-1, \mathbb{Z})$ and that the rows of U should be relatively primes.

3.2 Submanifolds and the kernel of G

In the following we would like to examine the structure of the kernel $\ker G$ of the metric G_{ij} on the commuting Killing vector fields $V_{(i)}$. As stated above, if we have a point q for which the kernel $\ker G$ is non-trivial then $\det G = 0$ at q . We define therefore the set

$$B = \{q \in \mathcal{N}_n \mid \det G(q) = 0\} \quad (3.4)$$

This is a codimension one hypersurface in $\mathcal{N}_n$, *i.e.* it is an $(n-1)$ -dimensional submanifold. As we shall see in the following this can be seen as part of the boundary of the manifold $\mathcal{N}_n$. Note that B need not be a connected set (see examples in Section 6). In the canonical form for the metric (2.10) B is the set of points with $r = 0$. In this coordinate system we can naturally equip B with the metric²

$$ds_B^2 = \Lambda_{\alpha\beta}|_{r=0} dz^\alpha dz^\beta \quad (3.5)$$

In the following we discuss for a given point $q \in B$ the Killing vector fields $v \in \ker G(q)$. We distinguish between Killing vector fields v which are space-like (time-like) near q , meaning that there exists a neighborhood $\mathcal{O}$ of q with respect to the manifold $\mathcal{N}_n$ such that $v^2 > 0$ ($v^2 < 0$) for any point in $\mathcal{O} - B$.

Define now the sets

$$Q_k = \{q \in \mathcal{N}_n \mid \dim \ker G(q) \geq k\} \quad (3.6)$$

for $k \in \{0, 1, \dots, p\}$. Clearly $Q_0 = \mathcal{N}_n$ and $Q_1 = B$. We examine now the properties of the sets Q_k .

²Note that for space-times solving the vacuum Einstein equations and obeying (2.6) this metric has determinant $\lambda = 1$.

Theorem 3.1 Consider a point $q \in Q_k - Q_{k+1}$. Then Q_k is a codimension k submanifold of $\mathcal{N}_n$ in a neighborhood of q .

Proof: Since $q \in Q_k - Q_{k+1}$ we have k linearly independent Killing vector fields $W_{(i)} \in \ker G$, $i = 1, 2, \dots, k$. We can always assume that at most one of these Killing vectors are time-like near q . If there were two of them which are time-like near q it follows from the fact that $V_{(1)}, \dots, V_{(p-1)}$ are space-like everywhere outside B that we can form a linear combination of the two Killing vectors which is space-like near q .

Assume first that all of these Killing vector fields are space-like near q . Then in order to avoid a conical singularity each of these Killing vector fields should generate a $U(1)$ isometry (see Section 3.1 for conditions on this, in particular one can infer from here that the k Killing vector fields are everywhere space-like). We can now find Riemannian Normal Coordinates $n^0, n^1, \dots, n^{D-1}$ in a neighborhood of q such that at q the D -dimensional metric is $ds^2 = \eta_{\mu\nu} dn^\mu dn^\nu$, the first derivatives of the metric at q in this coordinate system are zero, and such that the k Killing vectors can be written as ³

$$W_{(i)} = n^{2i} \frac{\partial}{\partial n^{2i+1}} - n^{2i+1} \frac{\partial}{\partial n^{2i}}, \quad i = 1, 2, \dots, k \quad (3.7)$$

That is, $W_{(i)}$ is the rotational Killing vector field in the plane (n^{2i}, n^{2i+1}) . Consider now the radii $\rho_i \equiv \sqrt{(n^{2i})^2 + (n^{2i+1})^2}$, $i = 1, 2, \dots, k$. First we observe that putting any of the $\rho_i > 0$ we have $\dim \ker G < k$. Secondly, for any point in the neighborhood of q we see that if $\rho_i = 0$ for all $i = 1, 2, \dots, k$ then we are in Q_k . Therefore we have shown that $Q_k = \{q' \in \mathcal{N}_n | \rho_i(q') = 0 \forall i = 1, 2, \dots, k\}$ in a neighborhood of q . Thus, we conclude that Q_k is a codimension k submanifold of $\mathcal{N}_n$ in a neighborhood of q .

Assume now that $W_{(k)}$ is time-like near q while $W_{(1)}, \dots, W_{(k-1)}$ all are space-like near q . Then $W_{(k)}$ generates $\mathbb{R}$ while $W_{(i)}$ generate $U(1)$ for $i = 1, 2, \dots, k-1$. We can now find Riemannian Normal Coordinates $n^0, n^1, \dots, n^{D-1}$ in a neighborhood of q such that at q the D -dimensional metric is $ds^2 = \eta_{\mu\nu} dn^\mu dn^\nu$, the first derivatives of the metric at q in this coordinate system are zero, and such that the k Killing vectors can be written as

$$W_{(k)} = n^0 \frac{\partial}{\partial n^1} + n^1 \frac{\partial}{\partial n^0}, \quad W_{(i)} = n^{2i} \frac{\partial}{\partial n^{2i+1}} - n^{2i+1} \frac{\partial}{\partial n^{2i}}, \quad i = 1, 2, \dots, k-1 \quad (3.8)$$

As in the other case we have the radii $\rho_i \equiv \sqrt{(n^{2i})^2 + (n^{2i+1})^2}$, $i = 1, 2, \dots, k-1$. We see that $W_{(k)}$ is the time-like Killing vector field in the Rindler space given by the coordinates (n^0, n^1) . We can therefore define the distance from the horizon in Rindler space as $\rho_k = \sqrt{(n^1)^2 - (n^0)^2}$. Proceeding with the argument the same way as above we see that $Q_k = \{q' \in \mathcal{N}_n | \rho_i(q') = 0 \forall i = 1, 2, \dots, k\}$ in a neighborhood of q and hence that Q_k is a codimension k submanifold of $\mathcal{N}_n$ in a neighborhood of q . $\square$

From this theorem we can infer the following corollary:

Corollary 3.2 Consider a point $q \in Q_{k+1} - Q_{k+2}$. Then Q_{k+1} is a codimension one submanifold of Q_k in a neighborhood of q . $\square$

We conclude from the above analysis that the structure of $\ker G$ naturally give rise to a hierarchy of submanifolds Q_k .

³A similar construction appeared in [5].

3.3 Domains and their directions

Suppose now that we consider a point $q \in B - Q_2$. Let furthermore $D \subset B - Q_2$ be the maximally possible connected set containing q . Write the coordinates for the point q as $(r, z^\alpha) = (0, z_*^\alpha)$. Since $\dim \ker G(q) = 1$ we can find a Killing vector field $W \in \ker G(q)$.

Suppose W is a space-like Killing vector field near q . In the following we aim to show that the linear space $\ker G$ is constant over D , *i.e.* that $W \in \ker G(q')$ for any point $q' \in D$.

One way to show the constancy of $\ker G$ over D is as follows [3]. We first observe that we can rigidly rotate G_{ij} such that $W = \partial/\partial x^1$. For $r \rightarrow 0$ and $z^\alpha \rightarrow z_*^\alpha$ we then have $G_{11} = c^2 r^2 + \mathcal{O}(r^3)$ with c a constant, and the entries of G_{ij} with $i, j \neq 1$ approaching a constant. In order for the space-time to be regular near q we need that $g_{rr} = e^{2(n-1)\nu}$ approaches a non-zero constant. We write this as $\nu \rightarrow c'$ for $r \rightarrow 0$ and $z^\alpha \rightarrow z_*^\alpha$. We also need that $A^1 \rightarrow 0$ for $r \rightarrow 0$ and $z^\alpha \rightarrow z_*^\alpha$. The metric (2.10) thus approaches

$$ds^2 = c^2 r^2 (dx^1)^2 + e^{2(n-1)c'} dr^2 + \sum_{i,j \neq 1} G_{ij}|_q (dx^i + A^i)(dx^j + A^j) + e^{2c'} \Lambda_{\alpha\beta}|_q dz^\alpha dz^\beta \quad (3.9)$$

for $r \rightarrow 0$ and $z^\alpha \rightarrow z_*^\alpha$. Consider now the R_{ij} part of the Ricci tensor (A.1). From requiring regularity we have that $R_{ij} = V_{(i}^\mu V_{j)}^\nu R_{\mu\nu}$ should be finite for $r \rightarrow 0$. Examining now R_{ii} for $i \neq 1$ one finds that $\tilde{g}^{ab} \partial_a G_{1i} \partial_b G_{1i} \rightarrow 0$ for $r \rightarrow 0$ and $z^\alpha \rightarrow z_*^\alpha$. This gives that $\partial_a G_{1i} = 0$ in q . Picking now any other point in D we can do the same. Since D is connected this means that $W = \partial/\partial x^1$ is in $\ker G$ everywhere in D . Undoing the rigid rotation we have shown that if $W \in \ker G(q)$ then $W \in \ker G(q')$ for any point $q' \in D$.

Another way to show the constancy of $\ker G$ over D is to employ the fact that to have a regular metric at q we need that W generates a $U(1)$ isometry [5]. Otherwise we get a conical singularity. This means we have

$$W = \sum_{i=1}^{p-1} q_i V_{(i)} \quad (3.10)$$

where the q_i 's are rational numbers. Since the norm of W is not significant we can choose to restrict the q_i 's to be relatively prime numbers. Now, for every point of $q' \in D$ we have an eigenvector $W_{q'} \in \ker G(q')$. But since $W_{q'}$ should vary continuously over D we see that one necessarily must have that $W_{q'} = W$. Thus we can conclude that $W \in \ker G$ everywhere in D hence $\ker G$ is constant over D .

Suppose instead W is a time-like Killing vector field near q . Making a rigid rotation of G_{ij} we can put $W = \partial/\partial x^0$. For the space-time to be regular near p the metric should approach

$$ds^2 = -c^2 r^2 (dx^0)^2 + e^{2(n-1)c'} dr^2 + \sum_{i,j \neq 0} G_{ij}|_q (dx^i + A^i)(dx^j + A^j) + e^{2c'} \Lambda_{\alpha\beta}|_q dz^\alpha dz^\beta \quad (3.11)$$

for $r \rightarrow 0$ and $z^\alpha \rightarrow z_*^\alpha$ where c and c' are constants. Just as above one can examine R_{ii} for $i \neq 0$ using (A.1) and find that $\partial_a G_{0i} = 0$ in p . Since p is any point in D this means that $W = \partial/\partial x^0 \in \ker G$ everywhere in D .

It follows furthermore from the fact that $W = \partial/\partial x^0 \in \ker G$ everywhere in D and the near- q metric (3.11) that for $r \rightarrow 0$ and $z^\alpha \rightarrow z_*^\alpha$ with $(0, z_*^\alpha) \in D$ the metric approaches

a Rindler space-time times a regular space of Euclidean signature. Therefore D is a Killing horizon of the Killing vector field $W = \partial/\partial x^0$.

We have thus shown above that the vectors in $\ker G$ are constant in the connected pieces of $B - Q_2$. With this, we can make the following definition:

Definition 3.3 *Let $q \in B - Q_2$ and let $W \in \ker G(q)$. A domain D containing q is the maximal connected set in B such that $q \in D$ and such that for any point $q' \in D$ we have $W \in \ker G(q')$. $\square$*

We can now consider all the distinct domains of B , write them as $D_1, \dots, D_N$. Clearly we have $D_i \cap D_j \subset Q_2$. From Corollary 3.2 we have that Q_2 can be seen locally as a submanifold of B of codimension one. This means that for $q \in Q_2$ any neighborhood of q in B will contain points in $B - Q_2$. This shows that the domains of B contains all points in Q_2 . Thus we can write $B = D_1 \cup D_2 \cup \dots \cup D_N$.

We have now shown the following theorem:

Theorem 3.4 *Let $D_1, \dots, D_N$ be the domains of B . We have that $B = D_1 \cup D_2 \cup \dots \cup D_N$. For each domain D_m we can find a Killing vector field W_m such that $W_m \in \ker G$ for all points in D_m . We call W_m the direction of the domain D_m . If W_m is space-like for $r \rightarrow 0$ we can write it in the form*

$$W_m = \sum_{i=1}^{p-1} q_i V_{(i)} \quad (3.12)$$

where the q_i 's are relatively prime numbers. Then W_m generates a $U(1)$ isometry and the generated flow has period 2π . In this case we say that the direction W_m is space-like.

If W_m is time-like for $r \rightarrow 0$ we can write it in the form

$$W_m = V_{(0)} + \sum_{i=1}^{p-1} \Omega_i V_{(i)} \quad (3.13)$$

and the domain D_m is a Killing horizon for the Killing vector field W_m . In this case we say that the direction W_m is time-like. $\square$

From Theorem 3.4 we can now define the domain structure of a solution:

Definition 3.5 *The domain structure of a solution is defined as the split-up of B in domains $B = D_1 \cup D_2 \cup \dots \cup D_N$ up to volume preserving diffeomorphisms, along with the directions W_m , $m = 1, 2, \dots, N$, of the domains.*

Our results above show that the domain structure of a given solution gives invariants of the solution (up to rigid transformations of the Killing vector fields as discussed in Section 3.1). In particular we have shown in Section 2.3 that the only left over coordinate transformations in the (r, z^α) coordinates are volume-preserving diffeomorphisms of the z^α coordinates.

Since the domain structure of a solution gives invariants of the solution it can characterize the solution. That is, it gives invariants that can help in distinguishing different solutions,

and it can furthermore provide information about the nature of the difference. A particular set of invariants derived from the domain structure consists of the volumes of the domains with respect to the metric (3.5). We call these invariants geometrical since they define in a coordinate-invariant way sizes of well-defined regions of the space-time as measured with the metric of the space-time.

In Section 7 we discuss for which type of solutions we can expect the domain structure to give a full characterization. We conjecture a uniqueness theorem for this type of solutions. We also discuss what extra information one has to add beyond the domain structure to give a full characterization of solutions coupled to gauge fields.

3.4 Reduction to the rod structure for $n = 2$

We consider here the special case $n = 2$, *i.e.* with $p = D - 2$ Killing vector fields. This is the case studied in [3, 5, 6].

Solutions of vacuum Einstein equations

We consider first solutions of vacuum Einstein equations with $p = D - 2$ commuting Killing vector fields. By theorem A.3 this means (under mild assumptions) that we can put the metric in the canonical form (2.9) which in this case reduces to

$$ds^2 = G_{ij}dx^i dx^j + e^{2\nu}(dr^2 + dz^2), \quad r^2 = |\det G_{ij}| \quad (3.14)$$

which is the canonical form of the metric found in [3]. Assuming $\mathcal{N}_2$ is simply connected we have $B = \mathbb{R}$, *i.e.* it is the z -axis for $r = 0$. Let $D_1, \dots, D_N$ be the domains of B with directions $W_1, \dots, W_N$. Then each domain corresponds to an interval of the z -axis. Thus, in the nomenclature of [3] each domain corresponds to a rod. Furthermore the direction of the rod is simply the direction of the domain. We also see that the volume preserving diffeomorphisms mentioned in Definition 3.5 here simply are the translations. The fact that the volumes of the domains are invariants corresponds to the statement that the lengths of the rods are invariants.

We thus regain the rod structure of [3]. We found moreover that the directions of the space-like rods can be written as (3.12) with the q_i being relatively prime numbers. This constraint has previously been found in [5].

General case

For the more general case of asymptotically flat or asymptotically Kaluza-Klein space solutions with $n = 2$ we get from (2.10) that the canonical form of the metric is

$$ds^2 = G_{ij}(dx^i + A^i)(dx^j + A^j) + e^{2\nu}(dr^2 + \lambda^2 dz^2), \quad r^2 = |\det G_{ij}| \quad (3.15)$$

$$\lambda \rightarrow 1 \text{ for } r \rightarrow \infty$$

This is more general than the form found in [3, 5, 6] since here we are not specific on what kinds of matter fields appear in the solution. Other than that the domains/rods are again

intervals on the z -axis defined by $r = 0$. The lengths of the domains/rods are measured by the metric

$$ds^2 = \lambda^2|_{r=0} dz^2 \quad (3.16)$$

These lengths are invariants of the black hole space-time. The domain/rod structure is defined up to translations. We have thus defined the rod structure for any asymptotically flat or asymptotically Kaluza-Klein black hole space-time with $p = D - 2$ commuting Killing vector fields. We can furthermore extend the rod structure to include non-asymptotically flat solutions [9].

4 Domain structure of six dimensional black holes

In this section we analyze the known asymptotically flat six-dimensional exact solutions of the vacuum Einstein equations. These are the Minkowski space, the Schwarzschild-Tangherlini black hole and the Myers-Perry black hole. They all have three Killing vector fields, which is the maximally possible number in six dimensions. In addition the Killing vector fields obey the condition (2.6). This means that the metrics can be put in the canonical form (2.9) with $p = n = 3$.

Minkowski space

The metric of six-dimensional Minkowski space is

$$ds^2 = -dt^2 + d\rho^2 + \rho^2(\mu_1^2 d\phi_1^2 + \mu_2^2 d\phi_2^2 + d\theta^2 + \cos^2 \theta d\psi^2) \quad (4.1)$$

with

$$\mu_1 = \sin \theta, \quad \mu_2 = \cos \theta \sin \psi, \quad \mu_3 = \cos \theta \cos \psi \quad (4.2)$$

and with the coordinate ranges $0 \leq \theta \leq \pi/2$ and $0 \leq \psi \leq \pi$. From (2.9) we see

$$r = \rho^2 \mu_1 \mu_2 = \rho^2 \sin \theta \cos \theta \sin \psi = \frac{1}{2} \rho^2 \sin(2\theta) \sin \psi \quad (4.3)$$

Using this we get $e^{-4\nu} = \rho^2 [\sin^2 \psi + \sin^2 \theta \cos^2 \psi]$. In order to fit in the canonical form (2.9) we need to find the z^α coordinates such that $g_{rz^\alpha} = 0$ and $\lambda = 1$. We make the following ansatz $z^\alpha = \rho^{k_\alpha} F_\alpha(\theta)(\cos \psi)^{l_\alpha}$, $\alpha = 1, 2$. Demanding that $g_{rz^\alpha} = 0$ gives that the functions $F_\alpha(\theta)$ are of the form $F_\alpha(\theta) = C_\alpha (\cos \theta)^{l_\alpha} (\cos 2\theta)^{\frac{k_\alpha - l_\alpha}{2}}$ where C_α are constants. One can furthermore infer that $\lambda = 1$ provided $C_1 C_2 = \pm 1 / (k_1 l_2 - k_2 l_1)$, $k_1 + k_2 = 3$ and $l_1 + l_2 = 1$. We choose therefore the coordinates

$$r = \frac{1}{2} \rho^2 \sin 2\theta \sin \psi, \quad z^1 = \rho \cos \theta \cos \psi, \quad z^2 = \frac{1}{2} \rho^2 \cos 2\theta \quad (4.4)$$

With this choice of coordinates the 6D flat space metric is put in the form (2.9).

We now analyze the domain structure of six-dimensional Minkowski space using the coordinates (4.4). This can be done by analyzing the coordinates z^α when $r = 0$. We find the

domain structure

$$\begin{aligned} W_1 &= \frac{\partial}{\partial \phi_1} , \quad D_1 = \{(z^1, z^2) \in \mathbb{R}^2 \mid z^2 \geq \frac{1}{2}(z^1)^2\} \\ W_2 &= \frac{\partial}{\partial \phi_2} , \quad D_2 = \{(z^1, z^2) \in \mathbb{R}^2 \mid z^2 \leq \frac{1}{2}(z^1)^2\} \end{aligned} \quad (4.5)$$

We see that $D_1 \cup D_2 = \mathbb{R}^2$. This domain structure is depicted in the top left diagram of Figure 1. We note that in terms of the (ρ, θ, ψ) coordinates the two domains correspond to $D_1 : \theta = 0$ and $D_2 : \psi = 0, \pi$.

Building on our parametrization of six-dimensional Minkowski space (4.4) we can now describe the boundary conditions that we wish to impose on six-dimensional asymptotically flat space-times. We consider here solutions with $p = 3$ such that one can write them in the form (2.10) in terms of coordinates (r, z^1, z^2) . We define the asymptotic region in (r, z^1, z^2) coordinates as $L \rightarrow \infty$ with $\sqrt{r}/L$, $(z^1)^2/L$ and z^2/L finite or going to zero where $L \equiv r + (z^1)^2 + |z^2|$. In this asymptotic region we require that the metric should asymptote to six-dimensional Minkowski space. This means in particular that we require the (r, z^1, z^2) coordinates to asymptote to Eq. (4.4). For the domain structure at $r = 0$ this means that for $(z^1)^2 + |z^2| \rightarrow \infty$ we have the two domains (4.5), with the border between the domains being at the curve $z^2 = (z^1)^2/2$ up to corrections of order $((z^1)^2 + |z^2|)^{-1/2}$.

Schwarzschild-Tangherlini black hole

The 6D Schwarzschild-Tangherlini black hole has the metric

$$ds^2 = -f dt^2 + \frac{d\rho^2}{f} + \rho^2(\mu_1^2 d\phi_1^2 + \mu_2^2 d\phi_2^2 + d\theta^2 + \cos^2 \theta d\psi^2) , \quad f = 1 - \frac{\rho_0^3}{\rho^3} \quad (4.6)$$

with the director cosines given by (4.2). We have from (2.9)

$$r = \rho^2 \sqrt{f} \mu_1 \mu_2 = \rho^2 \sqrt{f} \sin \theta \cos \theta \sin \psi = \frac{1}{2} \rho^2 \sqrt{f} \sin(2\theta) \sin \psi \quad (4.7)$$

From this one can easily compute $\exp(-4\nu)$ as function of (ρ, θ, ψ) . We need to impose that $g_{rz^\alpha} = 0$ and $\lambda = 1$. Make now the ansatz $z^1 = b_1(\rho) \cos \theta \cos \psi$ and $z^2 = b_2(\rho) \cos 2\theta$. Then $g_{rz^\alpha} = 0$ is equivalent to $2b'_1/b_1 = b'_2/b_2 = 8\rho^3/(4\rho^4 - \rho\rho_0^3)$. We therefore get the z^α coordinates

$$z^1 = \rho \left(1 - \frac{\rho_0^3}{4\rho^3}\right)^{\frac{1}{3}} \cos \theta \cos \psi , \quad z^2 = \frac{1}{2} \rho^2 \left(1 - \frac{\rho_0^3}{4\rho^3}\right)^{\frac{2}{3}} \cos 2\theta \quad (4.8)$$

Comparing this with (4.4) for six-dimensional Minkowski space we see that we have the right asymptotic behavior, as discussed above. One can also compute that $\lambda = 1$ which indeed is guaranteed by Eq. (2.8) and by choosing the right asymptotics.

The domain structure for the six-dimensional Schwarzschild-Tangherlini black hole as found from the coordinates (4.7) and (4.8) is given by

$$\begin{aligned} W_1 &= \frac{\partial}{\partial \phi_1} , \quad D_1 = \{(z^1, z^2) \in \mathbb{R}^2 \mid z^2 \geq K , \quad z^2 \geq \frac{1}{2}(z^1)^2\} \\ W_2 &= \frac{\partial}{\partial \phi_2} , \quad D_2 = \{(z^1, z^2) \in \mathbb{R}^2 \mid z^2 \leq (z^1)^2 - K , \quad z^2 \leq \frac{1}{2}(z^1)^2\} \\ W_3 &= \frac{\partial}{\partial t} , \quad D_3 = \{(z^1, z^2) \in \mathbb{R}^2 \mid (z^1)^2 - K \leq z^2 \leq K\} \end{aligned} \quad (4.9)$$

where we defined the constant $K \equiv (\rho_0^2/2)(3/4)^{2/3}$. This domain structure is depicted in the middle left diagram of Figure 1. We note that in terms of the (ρ, θ, ψ) coordinates the three domains correspond to $D_1 : \theta = 0$, $D_2 : \psi = 0, \pi$ and $D_3 : \rho = \rho_0$.

Myers-Perry black hole

The six-dimensional Myers-Perry black hole solution is [10]

$$ds^2 = -dt^2 + \sum_{i=1}^2 (\rho^2 + a_i^2) (d\mu_i^2 + \mu_i^2 d\phi_i^2) + \rho^2 d\mu_3^2 + \frac{\rho_0^3 \rho}{\Pi F} \left(dt - \sum_{i=1}^2 a_i \mu_i^2 d\phi_i \right)^2 + \frac{\Pi F d\rho^2}{\Pi - \rho \rho_0^3} \quad (4.10)$$

Here the director cosines are given by (4.2) and we have

$$F(\rho, \mu_i) = 1 - \sum_{i=1}^2 \frac{a_i^2 \mu_i^2}{\rho^2 + a_i^2}, \quad \Pi(\rho) = \prod_{i=1}^2 (\rho^2 + a_i^2) \quad (4.11)$$

The horizon is placed at $\rho = \rho_h$ which is defined as the largest real root of the equation $\Pi(\rho) = \rho \rho_0^3$. We find from (2.9)

$$r = \sqrt{\Pi - \rho \rho_0^3} \mu_1 \mu_2 = \frac{1}{2} \sqrt{\Pi - \rho \rho_0^3} \sin(2\theta) \sin \psi \quad (4.12)$$

We make the ansatz

$$z^1 = b_1(\rho) \cos \theta \cos \psi, \quad z^2 = b_2(\rho) \cos 2\theta + p(\rho) \cos^2 \theta \cos^2 \psi \quad (4.13)$$

Demanding $g_{rz^\alpha} = 0$ is equivalent to the equations

$$\frac{b'_1}{b_1} = \frac{4\rho^2 + 2(a_1^2 + a_2^2)}{4\rho^3 + 2\rho(a_1^2 + a_2^2) - \rho_0^3}, \quad \frac{b'_2}{b_2} = \frac{8\rho^2}{4\rho^3 + 2\rho(a_1^2 + a_2^2) - \rho_0^3} \quad (4.14)$$

$$(4\rho^3 + 2\rho(a_1^2 + a_2^2) - \rho_0^3)p' - 4(2\rho^2 + a_1^2 + a_2^2)p = 4a_2^2 b_2$$

Imposing the boundary conditions for $\rho \rightarrow \infty$ we get

$$b_1(\rho) = \rho \exp \left\{ - \int_{\rho/\rho_0}^{\infty} \frac{dx}{x(4x^3 + 2xA^2 - 1)} \right\} \quad (4.15)$$

$$b_2(\rho) = \frac{1}{2} \rho^2 \exp \left\{ - \int_{\rho/\rho_0}^{\infty} \frac{(2 - 4xA^2) dx}{x(4x^3 + 2xA^2 - 1)} \right\}$$

with $A^2 \equiv (a_1^2 + a_2^2)/\rho_0^2$. Considering the last equation in (4.14) we see that this is solved by

$$p = \frac{a_2^2}{a_1^2 + a_2^2} (b_1^2 - 2b_2) \quad (4.16)$$

where we fixed an integration constant by imposing the boundary condition $p(\rho)/b_2(\rho) \rightarrow 0$ for $\rho \rightarrow \infty$.

Comparing (4.13), (4.15) and (4.16) with (4.4) for six-dimensional Minkowski space we see that we have the right asymptotic behavior, as discussed above. One can compute that $\lambda = 1$ which again is guaranteed by Eq. (2.8) and by choosing the right asymptotics.

For the six-dimensional Myers-Perry black hole we find a domain structure with three domains D_1 , D_2 and D_3 with corresponding directions

$$W_1 = \frac{\partial}{\partial \phi_1}, \quad W_2 = \frac{\partial}{\partial \phi_2}, \quad W_3 = \frac{\partial}{\partial t} + \Omega_1 \frac{\partial}{\partial \phi_1} + \Omega_2 \frac{\partial}{\partial \phi_2} \quad (4.17)$$

We see that while the two first directions correspond to the two rotational Killing vector fields the third direction is instead the null Killing vector of the event horizon with the angular velocities given by $\Omega_i = a_i/(a_i^2 + r_h^2)$. The three domains are

$$\begin{aligned} D_1 &= \left\{ (z^1, z^2) \in \mathbb{R}^2 \left| z^1 = b_1(\rho)x, \quad z^2 = b_2(\rho) + p(\rho)x^2, \quad \rho \geq \rho_h, \quad |x| \leq 1 \right. \right\} \\ D_2 &= \left\{ (z^1, z^2) \in \mathbb{R}^2 \left| z^1 = b_1(\rho)y, \quad z^2 = b_2(\rho)(2y^2 - 1) + p(\rho)y^2, \quad \rho \geq \rho_h, \quad |y| \leq 1 \right. \right\} \\ D_3 &= \left\{ (z^1, z^2) \in \mathbb{R}^2 \left| z^1 = b_1(\rho_h)xy, \quad z^2 = b_2(\rho_h)(2y^2 - 1) + p(\rho_h)x^2y^2, \quad |x| \leq 1, \quad 0 \leq y \leq 1 \right. \right\} \end{aligned} \quad (4.18)$$

This domain structure is depicted in the bottom left diagram of Figure 1. We note that in terms of the (ρ, θ, ψ) coordinates the three domains correspond to $D_1 : \theta = 0$, $D_2 : \psi = 0, \pi$ and $D_3 : \rho = \rho_h$.

5 Domain structure of seven dimensional black holes

In this section we analyze the known asymptotically flat seven-dimensional exact solutions of the vacuum Einstein equations. These are the Minkowski space, the Schwarzschild-Tangherlini black hole and the Myers-Perry black hole. They all have four Killing vector fields, which is the maximally possible number in seven dimensions. In addition the Killing vector fields obey the condition (2.6). This means that the metrics can be put in the canonical form (2.9) with $p = 4$ and $n = 3$.

Minkowski space

The metric of seven-dimensional Minkowski space is

$$ds^2 = -dt^2 + d\rho^2 + \rho^2(\mu_1^2 d\phi_1^2 + \mu_2^2 d\phi_2^2 + \mu_3^2 d\phi_3^2 + d\theta^2 + \cos^2 \theta d\psi^2) \quad (5.1)$$

with the director cosines

$$\mu_1 = \sin \theta, \quad \mu_2 = \cos \theta \sin \psi, \quad \mu_3 = \cos \theta \cos \psi \quad (5.2)$$

and with the coordinate ranges $0 \leq \theta, \psi \leq \pi/2$. Using (2.9) we see that

$$r = \rho^3 \mu_1 \mu_2 \mu_3 = \rho^3 \sin \theta \cos^2 \theta \sin \psi \cos \psi = \frac{1}{2} \rho^3 \sin \theta \cos^2 \theta \sin 2\psi \quad (5.3)$$

From this we get $e^{-4\nu} = \rho^4 \cos^2 \theta [4 \sin^2 \theta + \cos^2 \theta \sin^2(2\psi)]/4$. In order to fit in the canonical form (2.9) we need to find the z^α coordinates such that $g_{rz^\alpha} = 0$ and $\lambda = 1$. We make the following ansatz $z^\alpha = \rho^{k_\alpha} F_\alpha(\theta)(\cos 2\psi)^{l_\alpha}$ with $\alpha = 1, 2$. Demanding that $g_{rz^\alpha} = 0$ gives

that the functions $F_\alpha(\theta)$ are of the form $F_\alpha(\theta) = C_\alpha(\cos \theta)^{2l_\alpha}(3\cos^2 \theta - 2)^{\frac{k_\alpha}{2}-l_\alpha}$ where C_α are constants. One can furthermore infer that $\lambda = 1$ provided $4C_1C_2 = \pm 1/(l_1k_2 - k_1l_2)$, $k_1 + k_2 = 4$ and $l_1 + l_2 = 1$. We choose therefore the z^α coordinates

$$z^1 = \frac{1}{2}\rho^2 \cos^2 \theta \cos 2\psi, \quad z^2 = \frac{1}{4}\rho^2(3\cos^2 \theta - 2) \quad (5.4)$$

We now consider the domain structure of seven-dimensional Minkowski space using the coordinates (5.3) and (5.4). This can be done by analyzing the coordinates z^α when $r = 0$. We find the domain structure

$$\begin{aligned} W_1 &= \frac{\partial}{\partial \phi_1}, \quad D_1 = \left\{ (z^1, z^2) \in \mathbb{R}^2 \left| z^2 \geq \frac{1}{2}|z^1| \right. \right\} \\ W_2 &= \frac{\partial}{\partial \phi_2}, \quad D_2 = \left\{ (z^1, z^2) \in \mathbb{R}^2 \left| z^1 \geq 0, z^2 \leq \frac{1}{2}z^1 \right. \right\} \\ W_3 &= \frac{\partial}{\partial \phi_3}, \quad D_3 = \left\{ (z^1, z^2) \in \mathbb{R}^2 \left| z^1 \leq 0, z^2 \leq -\frac{1}{2}z^1 \right. \right\} \end{aligned} \quad (5.5)$$

We see that $D_1 \cup D_2 \cup D_3 = \mathbb{R}^2$. This domain structure is depicted in the top right diagram of Figure 1. We note that in terms of the (ρ, θ, ψ) coordinates the three domains correspond to $D_1 : \theta = 0$, $D_2 : \psi = 0$ and $D_3 : \psi = \pi/2$. The origin of Minkowski space $\rho = 0$ is seen to be the common intersection point of all of the three domains. This makes sense since the origin is the only point which is a fixed point of rotation in all of the three rotation planes.

Building on our parametrization of seven-dimensional Minkowski space given by Eqs. (5.3) and (5.4) we can now describe the boundary conditions that we wish to impose on seven-dimensional asymptotically flat space-times. We consider here solutions with $p = 3$ such that one can write them in the form (2.10) in terms of coordinates (r, z^1, z^2) . We define the asymptotic region in (r, z^1, z^2) coordinates as $L \rightarrow \infty$ with $r^{2/3}/L$, z^1/L and z^2/L finite or going to zero where $L \equiv r^{2/3} + |z_1| + |z_2|$. In this asymptotic region we require that the metric should asymptote to seven-dimensional Minkowski space. This means in particular that we require the (r, z^1, z^2) coordinates to asymptote to Eqs. (5.3) and (5.4). For the domain structure at $r = 0$ this means that for $|z_1| + |z_2| \rightarrow \infty$ we have the three domains (5.5), with the border between the domains being at the curves $z^2 = |z^1|/2$ and $z^1 = 0$ for $z^2 \leq 0$ up to corrections of order $(|z_1| + |z_2|)^{-1}$.

Schwarzschild-Tangherlini black hole

The 7D Schwarzschild-Tangherlini black hole has the metric

$$ds^2 = -f dt^2 + \frac{d\rho^2}{f} + \rho^2(\mu_1^2 d\phi_1^2 + \mu_2^2 d\phi_2^2 + \mu_3^2 d\phi_3^2 + d\theta^2 + \cos^2 \theta d\psi^2), \quad f = 1 - \frac{\rho_0^4}{\rho^4} \quad (5.6)$$

with the director cosines given by (5.2). We get

$$r = \rho^3 \sqrt{f} \mu_1 \mu_2 \mu_3 = \rho^3 \sqrt{f} \sin \theta \cos^2 \theta \sin \psi \cos \psi = \frac{1}{2} \rho^3 \sqrt{f} \sin \theta \cos^2 \theta \sin 2\psi \quad (5.7)$$

From this one can easily compute $\exp(-4\nu)$ as function of (ρ, θ, ψ) . Make now the ansatz $z^1 = b_1(\rho) \cos^2 \theta \cos 2\psi$ and $z^2 = b_2(\rho)(3\cos^2 \theta - 2)$. Imposing $g_{rz^\alpha} = 0$ is equivalent to

$b'_1/b_1 = b'_2/b_2 = 6\rho^3/(3\rho^4 - \rho_0^4)$. We get therefore the z^α coordinates

$$z^1 = \frac{1}{2}\sqrt{\rho^4 - \frac{\rho_0^4}{3}} \cos^2 \theta \cos 2\psi, \quad z^2 = \frac{1}{4}\sqrt{\rho^4 - \frac{\rho_0^4}{3}} (3 \cos^2 \theta - 2) \quad (5.8)$$

Comparing this with (5.3) and (5.4) for seven-dimensional Minkowski space we see that we have the right asymptotic behavior, as discussed above. One can compute that $\lambda = 1$ which is guaranteed by Eq. (2.8) and by choosing the right asymptotics.

The domain structure for the seven-dimensional Schwarzschild-Tangherlini black hole as found from the coordinates (5.7) and (5.8) is given by

$$\begin{aligned} W_1 &= \frac{\partial}{\partial \phi_1}, \quad D_1 = \left\{ (z^1, z^2) \in \mathbb{R}^2 \left| z^2 \geq \frac{1}{2}|z^1|, z^2 \geq \frac{\rho_0^2}{2\sqrt{6}} \right. \right\} \\ W_2 &= \frac{\partial}{\partial \phi_2}, \quad D_2 = \left\{ (z^1, z^2) \in \mathbb{R}^2 \left| z^1 \geq 0, z^2 \leq \frac{1}{2}z^1, z^2 \leq \frac{3}{2}z^1 - \frac{\rho_0^2}{\sqrt{6}} \right. \right\} \\ W_3 &= \frac{\partial}{\partial \phi_3}, \quad D_3 = \left\{ (z^1, z^2) \in \mathbb{R}^2 \left| z^1 \leq 0, z^2 \leq -\frac{1}{2}z^1, z^2 \leq -\frac{3}{2}z^1 - \frac{\rho_0^2}{\sqrt{6}} \right. \right\} \\ W_4 &= \frac{\partial}{\partial t}, \quad D_4 = \left\{ (z^1, z^2) \in \mathbb{R}^2 \left| \frac{3}{2}|z^1| - \frac{\rho_0^2}{\sqrt{6}} \leq z^2 \leq \frac{\rho_0^2}{2\sqrt{6}} \right. \right\} \end{aligned} \quad (5.9)$$

This domain structure is depicted in the middle right diagram of Figure 1. We note that in terms of the (ρ, θ, ψ) coordinates the four domains correspond to $D_1 : \theta = 0$, $D_2 : \psi = 0$, $D_3 : \psi = \pi/2$ and $D_4 : \rho = \rho_0$.

Myers-Perry black hole

The seven-dimensional Myers-Perry black hole solution is [10]

$$ds^2 = -dt^2 + \sum_{i=1}^3 (\rho^2 + a_i^2) (d\mu_i^2 + \mu_i^2 d\phi_i^2) + \frac{\rho_0^{D-3} \rho^2}{\Pi F} \left(dt - \sum_{i=1}^3 a_i \mu_i^2 d\phi_i \right)^2 + \frac{\Pi F d\rho^2}{\Pi - \rho^2 \rho_0^4} \quad (5.10)$$

with the director cosines given by (5.2) and we have

$$F(\rho, \mu_i) = 1 - \sum_{i=1}^3 \frac{a_i^2 \mu_i^2}{\rho^2 + a_i^2}, \quad \Pi(\rho) = \prod_{i=1}^3 (\rho^2 + a_i^2) \quad (5.11)$$

The horizon is placed at $\rho = \rho_h$ which is defined as the largest real root of the equation $\Pi(\rho) = \rho^2 \rho_0^4$. From (2.9) we find

$$r = \sqrt{\Pi - \rho^2 \rho_0^4} \mu_1 \mu_2 \mu_3 = \frac{1}{2} \sqrt{\Pi - \rho^2 \rho_0^4} \sin \theta \cos^2 \theta \sin 2\psi \quad (5.12)$$

We use the following ansatz for z^α

$$z^\alpha = z_0^\alpha + p_\alpha(\rho) \cos^2 \theta \cos 2\psi + q_\alpha(\rho) (3 \cos^2 \theta - 2) \quad (5.13)$$

The orthogonality conditions $g_{rz^\alpha} = 0$ are equivalent to the relations

$$\begin{aligned} \rho(a_2^2 - a_3^2)p_\alpha &= -3\rho(2\rho^2 + a_2^2 + a_3^2)q_\alpha + (3\rho^4 + 2\rho^2 a^2 + B^4 - \rho_0^4)q'_\alpha \\ 3\rho(a_2^2 - a_3^2)q_\alpha &= -\rho(6\rho^2 + 4a_1^2 + a_2^2 + a_3^2)p_\alpha + (3\rho^4 + 2\rho^2 a^2 + B^4 - \rho_0^4)p'_\alpha \end{aligned} \quad (5.14)$$




Figure 1: On the left side are shown the domain structures for the six-dimensional Minkowski space (top left), the Schwarzschild-Tangherlini black hole (middle left) with $\rho_0^3 = 4/3$ and the Myers-Perry black hole (bottom left) with $\rho_0^3 = 4/3$, $a_1 = 1/4$ and $a_2 = 4/5$. On the right side are shown the domain structures for the seven-dimensional Minkowski space (top right), the Schwarzschild-Tangherlini black hole (middle right) with $\rho_0^4 = 6$ and the Myers-Perry black hole (bottom right) with $\rho_0^4 = 6$, $a_1 = 3/2$, $a_2 = 3/4$ and $a_3 = 1/3$.

where we defined for convenience $a^2 \equiv a_1^2 + a_2^2 + a_3^2$, $B^4 \equiv a_1^2 a_2^2 + a_1^2 a_3^2 + a_2^2 a_3^2$ and $C^4 \equiv a_1^4 + a_2^4 + a_3^4 - B^4$. From these relations one can infer that p_α and q_α solve the same second order ODE which has the two independent solutions

$$F_{\pm}(\rho) \equiv \sqrt{3\rho^4 + 2a^2\rho^2 + B^4 - \rho_0^4} \exp \left\{ \pm \frac{C^2}{\sqrt{3\rho_0^4 + C^4}} \operatorname{arctanh} \left(\frac{\sqrt{3\rho_0^4 + C^4}}{3\rho^2 + a^2} \right) \right\} \quad (5.15)$$

Write now

$$p_\alpha(\rho) = p_\alpha^+ F_+(\rho) + p_\alpha^- F_-(\rho), \quad q_\alpha(\rho) = q_\alpha^+ F_+(\rho) + q_\alpha^- F_-(\rho) \quad (5.16)$$

One set of constraints on $p_\alpha^\pm$ and $q_\alpha^\pm$ comes from demanding that z^1 and z^2 asymptotes to (5.4) for $\rho \rightarrow \infty$. This can be worked out using that $F_{\pm}(\rho) \simeq \sqrt{3}\rho^2$ for $\rho \rightarrow \infty$. Another set of constraints is that the equations (5.14) should be satisfied. This fixes

$$p_1^\pm = 2q_2^\mp = \mp \frac{2a_1^2 - a_2^2 - a_3^2 \mp 2C^2}{8\sqrt{3}C^2}, \quad q_1^\pm = \frac{2}{3}p_2^\pm = \mp \frac{a_2^2 - a_3^2}{8\sqrt{3}C^2} \quad (5.17)$$

We furthermore impose that $z^1 \rightarrow 0$ for $\rho \rightarrow \infty$ when $\theta = \pi/2$ and $z^2|_{\psi=0} + z^2|_{\psi=\pi/2} \rightarrow 0$ for $\rho \rightarrow \infty$ when $3\cos^2\theta = 2$. This fixes $z_0^1 = -(a_2^2 - a_3^2)/6$ and $z_0^2 = 0$.

Comparing the coordinates (5.12)-(5.13) with those of seven-dimensional Minkowski space (5.3) and (5.4) we see that we have the right asymptotic behavior, as discussed above. One can again compute that $\lambda = 1$ which is guaranteed by Eq. (2.8) and by choosing the right asymptotics.

For the seven-dimensional Myers-Perry black hole we find a domain structure with four domains D_1, D_2, D_3 and D_4 with corresponding directions

$$W_1 = \frac{\partial}{\partial\phi_1}, \quad W_2 = \frac{\partial}{\partial\phi_2}, \quad W_3 = \frac{\partial}{\partial\phi_3}, \quad W_4 = \frac{\partial}{\partial t} + \Omega_1 \frac{\partial}{\partial\phi_1} + \Omega_2 \frac{\partial}{\partial\phi_2} + \Omega_3 \frac{\partial}{\partial\phi_3} \quad (5.18)$$

We see that while the three first directions correspond to the three rotational Killing vector fields the fourth direction is instead the null Killing vector of the event horizon with the angular velocities given by $\Omega_i = a_i/(a_i^2 + r_h^2)$. The four domains are

$$\begin{aligned} D_1 &= \left\{ (z^1, z^2) \in \mathbb{R}^2 \left| z^\alpha = z_0^\alpha + p_\alpha(\rho)x + q_\alpha(\rho), \rho \geq \rho_h, |x| \leq 1 \right. \right\} \\ D_2 &= \left\{ (z^1, z^2) \in \mathbb{R}^2 \left| z^\alpha = z_0^\alpha + p_\alpha(\rho)y + q_\alpha(\rho)(3y - 2), \rho \geq \rho_h, 0 \leq y \leq 1 \right. \right\} \\ D_3 &= \left\{ (z^1, z^2) \in \mathbb{R}^2 \left| z^\alpha = z_0^\alpha - p_\alpha(\rho)y + q_\alpha(\rho)(3y - 2), \rho \geq \rho_h, 0 \leq y \leq 1 \right. \right\} \\ D_4 &= \left\{ (z^1, z^2) \in \mathbb{R}^2 \left| z^\alpha = z_0^\alpha + p_\alpha(\rho_h)yx + q_\alpha(\rho_h)(3y - 2), |x| \leq 1, 0 \leq y \leq 1 \right. \right\} \end{aligned} \quad (5.19)$$

This domain structure is depicted in the bottom right diagram of Figure 1. We note that in terms of the (ρ, θ, ψ) coordinates the four domains correspond to $D_1 : \theta = 0$, $D_2 : \psi = 0$, $D_3 : \psi = \pi/2$ and $D_4 : \rho = \rho_h$.

6 Possible new domain structures in six and seven dimensions

In this section we examine the possible domain structures one can have for asymptotically flat solutions in six and seven dimensions with $D - 3$ commuting linearly independent Killing vector fields. We illustrate the domain structure diagrams in a different fashion than in Sections 4 and 5 since here we do not care about all details of the domain structure.

6.1 Six-dimensional asymptotically flat space-times

We consider here the possible domain structures of asymptotically flat six-dimensional black hole space-times with three commuting linearly independent Killing vector fields.

In the first diagram of Figure 2 we have depicted the domain structure of six-dimensional Minkowski space. Here the upper domain has direction $\partial/\partial\phi_1$ and the lower domain direction $\partial/\partial\phi_2$. These two domains correspond to the set of fixed points of the rotations in two rotation planes of six-dimensional Minkowski space. The idea is now to examine all the possible ways in which we can put a domain with a time-like direction corresponding to an event horizon in this domain structure diagram. We represent the event horizon domain as a filled area.

Over this domain is fibred two circles parameterized by the two rotation angles ϕ_1 and ϕ_2 . The topology of the event horizon is now determined from where these two circles shrink to zero at the boundary of the domain. This give rise to three distinct types of event horizons corresponding to S^4 , $S^1 \times S^3$ or $S^2 \times S^2$ topology as we discuss below. Another possibility is that the domain structure do not live in the plane $\mathbb{R}^2$ but in a disconnected space. As we discuss below this can give rise to an event horizon with $T^2 \times S^2$ topology.




Figure 2: Domain structure for six-dimensional Minkowski space and four possible domain structures for six-dimensional asymptotically flat black holes with a single event horizon.

The first possibility is to put the event horizon domain across the boundary of the two rotational domains. This is depicted in the second diagram of Figure 2 where we chose for convenience the filled area to have a shape corresponding to an area in between the branches of a parabola. We see that the boundary of the domain is divided in two parts, one in which the first circle is shrunk to zero, the other in which the second circle is shrunk to zero. This corresponds to the topology of a four-sphere. This is shown explicitly in Appendix B. From comparing with Figure 1 we see that this domain structure indeed is equivalent to those of the six-dimensional Schwarzschild-Tangherlini and Myers-Perry black hole.

The second possibility is to put the event horizon domain away from the curve separating the two rotational domains. This gives two possibilities, depending on whether we put it above or below. However these two possibilities are equivalent by relabeling the two rotation planes. We have illustrated one of the possibilities in the third diagram of Figure 2. In such a space-time the event horizon is topologically an $S^1 \times S^3$. This is seen from the fact that we again have two circles fibred over a disc but on the boundary the one parameterized by ϕ_1 shrinks to zero while the other one is of non-zero size everywhere on the event horizon. As shown explicitly in Appendix B a circle fibred over a disc for which the circle shrinks to zero at the boundary of the disc corresponds to a three-sphere topology. Thus, the domain structure corresponds to a black ring in six dimensions. Approximate metrics for neutral black rings in the ultraspinning regime have been found in [11] and described using the Blackfold approach

in [12].

The third possibility is that the event horizon domain is disconnected from the rotational domains. This can happen if the event horizon is displaced from the fixed points of rotations in both of the rotation planes. In [12] an example of this called a black torus is described with $T^2 \times S^2$ topology using the Blackfold approach again in the ultra-spinning regime. This is realized by having the domain submanifold $B = \mathbb{R}^2 \cup S^2$. This is concretely realized as having the domain plane parameterized by (z^1, z^2) being multi-valued so that for the (z^1, z^2) values where we have the event horizon domain we have three sheets of the domain plane – one sheet corresponding to the domain structure of the six-dimensional Minkowski space and the two other sheets disconnected from this being the two sides of a two-sphere projected on to a plane, see Appendix B for an explicit parametrization of this. In the fourth diagram of Figure 2 we have depicted this domain structure where the dashed line represents that the event horizon domain is disconnected from the two rotational domains. Clearly a space-time with such a domain structure has an event horizon with $T^2 \times S^2$ topology, with $T^2 = S^1 \times S^1$ being a rectangular torus, since the two circles do not shrink to zero at any point on the event horizon domain.

Finally, the fourth possibility for a domain structure is depicted in the fifth diagram of Figure 2. We see that the event horizon here is shaped as a piece of a ring. The event horizon can be seen to have an $S^2 \times S^2$ topology since in the angular direction we have that the ϕ_2 circle shrinks to zero in the two ends, while in the radial direction the ϕ_1 circle shrinks to zero in the two ends. Unlike the three above domain structures we do not have any evidence that this domain structure corresponds to a regular black hole space-time. However, numerical evidence for a static black hole space-time with this domain structure, though with a conical singularity, has been found in [13]. In the Blackfold approach [12] this kind of event horizon topology has also been considered in the limit in which one sphere is much larger than the other. It was found that the sphere cannot be supported by a single large angular momentum in this limit. However, it is conceivable that the solution can be made regular by having two angular momenta turned on, one for each sphere.

Multiple horizons

It is interesting to consider the combinations one can make of the above domain structures. For simplicity we restrict ourselves to the first two possibilities depicted in Figure 2. First we can make a Black Saturn, i.e. a black ring with a black hole in the center. This corresponds to the domain structure depicted in the first diagram of Figure 3. In five dimensions such a solution has been found in [14]. We can also make a di-ring, i.e. two rings which are concentric and rotating in the same rotation plane. This corresponds to the domain structure of the second diagram of Figure 3. In five dimensions such a solution has been found in [15]. Finally we can imagine two bicycling black rings. These rotate in two orthogonal rotation planes. This corresponds to the domain structure of the third diagram of Figure 3. In five dimensions such a solution has been found in [16].




Figure 3: Three possible domain structures for six-dimensional asymptotically flat black holes with two separate event horizons.

6.2 Seven-dimensional asymptotically flat space-times

We consider here the possible domain structures of asymptotically flat seven-dimensional black hole space-times with four commuting linearly independent Killing vector fields.

In the first diagram of Figure 4 we have depicted seven-dimensional Minkowski space. Here the upper domain has direction $\partial/\partial\phi_1$, the right domain has direction $\partial/\partial\phi_2$ and the left domain has direction $\partial/\partial\phi_3$. These three domains correspond to the set of fixed points of the rotations in three rotation planes of seven-dimensional Minkowski space. We now want to examine all the possible ways in which we can put a domain with a time-like direction corresponding to an event horizon in this diagram. We represent this domain as a filled area. Over this domain is fibred three circles parameterized by the three rotation angles ϕ_1 , ϕ_2 and ϕ_3 . The topology of the event horizon is now determined from where these three circles shrink to zero at the boundary of the domain. This give rise to four distinct types of event horizons corresponding to the topologies S^5 , $S^1 \times S^4$, $T^2 \times S^3$ and $S^3 \times S^2$ as we discuss below. It is furthermore possible to draw a domain structure that gives rise to a topology with identifications of the five-sphere as we describe below. Another possibility is that the domain structure does not live in the plane $\mathbb{R}^2$ but in a disconnected space. As we discuss below this can give rise to an event horizon with $T^3 \times S^2$ topology.

The first possibility is that the event horizon domain covers the origin of the seven-dimensional Minkowski space - i.e. the point belonging to all three rotational domains. This is depicted in the second diagram of Figure 4 where the filled area for convenience has the shape of a triangle. We see that the boundary of the domain is divided in three parts, one for each side of the triangle. At each side of the triangle a different circle shrinks to zero. From this one can infer that the event horizon has topology of a five-sphere. This is shown explicitly in Appendix B. Comparing this domain structure with Figure 1 we see that it is equivalent to those of the seven-dimensional Schwarzschild-Tangherlini black hole and Myers-Perry black hole.

The second possibility is that the event horizon domain crosses one of the curves dividing the three rotational domains. This domain structure is depicted in the third diagram of Figure 4. It gives rise to an event horizon with $S^1 \times S^4$ topology where the S^1 corresponds to the ϕ_1 circle since that is finite everywhere on the event horizon domain. Instead with respect to the ϕ_2 and ϕ_3 circles we see that the boundary of the event horizon domain is divided in two parts, one part where the ϕ_2 circle shrinks to zero, the other where the ϕ_3 shrinks to zero.




Figure 4: Domain structure for seven-dimensional Minkowski space and five possible domain structures for seven-dimensional asymptotically flat black holes with a single event horizon.

As shown in Section 6.1 this corresponds to an S^4 topology. Thus, this domain structure corresponds to a black ring in seven dimensions. Approximate metrics for neutral black rings in the ultraspinning regime have been found in [11] and described using the Blackfold approach in [12].

The third possibility is that the event horizon domain do not cross any of the curves dividing the three rotational domains. This domain structure is depicted in the fourth diagram of Figure 4. This corresponds to an event horizon with $T^2 \times S^3$ topology where the rectangular torus $T^2 = S^1 \times S^1$ corresponds to the ϕ_2 and ϕ_3 circles since they do not shrink to zero on the event horizon domain. Instead the ϕ_1 circle shrinks to zero on the boundary of the event horizon domain hence this gives rise to the S^3 part of the topology, as shown in Section 6.1. Thus, this domain structure corresponds to a so-called black torus which has been described using the Blackfold approach in [12].

The fourth possibility is that the event horizon domain covers an area in between two of the curves dividing the three rotational domains. This domain structure is depicted in the fifth diagram of Figure 4 with the shape of a piece of a ring. This corresponds to an event horizon with $S^3 \times S^2$ topology. This is seen from the fact that the boundary of the domain is split up in four intervals, each corresponding to a side of the domain, according to which circle shrinks to zero. For the upper and lower sides the ϕ_1 circle shrinks to zero while for the left and right sides either the ϕ_2 or ϕ_3 circle shrinks to zero. This clearly gives an $S^3 \times S^2$ topology since when we go from boundary to boundary in the angular direction we go from shrinking the ϕ_2 circle to shrinking the ϕ_3 circle, thus giving a three-sphere, and when we go from boundary to boundary in the radial direction we shrink the ϕ_1 circle at both boundaries thus giving a two-sphere. In the Blackfold approach of [12] such an event horizon topology has been found in the limit where the S^2 is much smaller than the S^3 , corresponding to the

limit where the upper and lower sides in the fifth diagram of Figure 4 are very close.

The domain structure giving a $S^3 \times S^2$ topology is of particular interest since we see that it has a finite size domain with a space-like direction. This is very reminiscent of the rod structure of the five-dimensional black ring where one has a finite space-like rod. In particular this means that the domain structure has two invariants corresponding to the areas of the two domains of finite size, as measured using the metric (3.5).

The finite size space-like domain also provide a possible generalization. If we let the direction of this finite size domain be

$$W = \frac{\partial}{\partial \phi_1} + q \frac{\partial}{\partial \phi_2} \quad (6.1)$$

with q an integer, then we see that we have a Lens space $L(q, 1) = S^3/\mathbb{Z}_q$ when going in the radial direction. Instead in the angular direction we still have an S^3 in terms of the ϕ_2 and ϕ_3 circles. Thus, the topology of the event horizon is now $S^5/\mathbb{Z}_q$, which is a five-dimensional Lens space. This is reminiscent of what happens for five-dimensional black holes where one can get a three-dimensional Lens space by changing the direction of the finite space-like rod in the rod structure of the black ring [5, 17].

Finally, the last possibility considered here is that the event horizon domain is disconnected from the rotational domains. Thus the event horizon is displaced from the fixed points of rotation in all three rotation planes. This works the same way as in six dimensions. The domain submanifold is $B = \mathbb{R}^2 \cup S^2$ and it can again be viewed as a three-sheeted plane. This gives a three-torus topology $T^3 \times S^2$, with $T^3 = S^1 \times S^1 \times S^1$ a rectangular three-torus, and such a black three-torus have indeed been described by the Blackfold approach in [12]. We depicted the domain structure in the sixth diagram of Figure 4.

Multiple horizons

Just as in six dimensions it is again interesting to consider the combinations one can make of the above examples of domain structures for seven-dimensional black holes. Examples of this include the seven-dimensional version of the Black Saturn, see first diagram of Figure 5, a black hole (S^5 topology) with a black torus ($T^2 \times S^3$ topology) around, see second diagram of Figure 5, a black ring ($S^1 \times S^4$ topology) with a black torus ($T^2 \times S^3$ topology) around, see third diagram of Figure 5, and an black hole (S^5 topology) with a black three-sphere around ($S^3 \times S^2$ topology), see fourth diagram of Figure 5.




Figure 5: Four possible domain structures for seven-dimensional asymptotically flat black holes with two separate event horizons.

7 Discussion and outlook

In this paper we have introduced the domain structure for black hole space-times. We have shown that the domain structure provides invariants for a given space-time and that these invariants therefore can be part of the characterization of the space-time. A natural question following this is whether these invariants are enough to give a complete characterization of a black hole space-time.

We first restrict ourselves to solutions of the vacuum Einstein equations, and assume furthermore that the orthogonality condition (2.6) is obeyed. For stationary and asymptotically flat solutions with $[(D-1)/2]$ rotational Killing vector fields we have the highest number of Killing vector fields possible for solutions which are not Minkowski space. It is natural to make the following conjecture:⁴

Conjecture 7.1 *Let two D -dimensional regular and stationary asymptotically flat solutions of the vacuum Einstein equations be given, both with a single connected event horizon and with $[(D-1)/2]$ commuting rotational Killing vector fields obeying the orthogonality condition (2.6). Let the two solutions have the same mass and angular momenta. Assume that the set B is connected for both solutions. Then the two solutions are the same if and only if they have the same domain structure. $\square$*

For $D = 5$ this is shown to be true [5] following the uniqueness hypothesis of [3]. However, for $D > 5$ it is clear that one cannot apply the techniques used for $D = 4, 5$. The problem is that the metric $\tilde{g}_{ab}$ on $\mathcal{N}_n$ is not decoupled from the Killing vector metric G_{ij} in the Einstein equations. Thus, when given two solutions with the same domain structure they will, generically, have both two different G_{ij} metrics as well as two different $\tilde{g}_{ab}$ metrics. This means proving a uniqueness theorem is a considerably more challenging task than for $D = 5$ where it was enough to generalize the methods introduced for $D = 4$ [18, 5].

Consider now the general case, *i.e.* without the orthogonality condition (2.6) or restrictions on what matter fields are present. Here we observe that the domain structure in general is not enough to fully characterize a solutions. To see this we can use a lesson from the paper [19] where a ring with a dipole charge was found. This gives infinite non-uniqueness of the solution since the ring carries no net charge, as measured at infinity. Clearly the domain structure cannot carry information on the dipole charge thus it is evident that one needs to supplement the domain structure invariants with information about locally measured charges, such as the dipole charge (see [20] for a general exposition on dipole charges and other local charges). It would be very interesting to pursue this problem further to find a general way to specify the dipole charges - as well as other types of local charges - for the event horizon domains. Combined with the domain structure this could lead to a full characterization of asymptotically flat black hole space-times with $[(D-1)/2]$ rotational Killing vector fields.

Another direction which is interesting to consider is asymptotically flat solutions with less than $[(D-1)/2]$ rotational Killing vector fields. As an example we can take the case of $D = 5$.

⁴Note that we added an assumption on connectedness of B since it is unclear whether the domain structure contains enough information to parameterize a situation with disconnected B .

Consider a stationary, but non-static, black hole. Write the null Killing vector of the event horizon as

$$W = \frac{\partial}{\partial t} + \Omega_1 \frac{\partial}{\partial \phi_1} + \Omega_2 \frac{\partial}{\partial \phi_2} \quad (7.1)$$

Then we know from the Rigidity theorems of [8] that W is a Killing vector field of the space-time. Thus, the space-time have the two Killing vector fields

$$V_{(0)} = \frac{\partial}{\partial t}, \quad V_{(1)} = p \frac{\partial}{\partial \phi_1} + q \frac{\partial}{\partial \phi_2} \quad (7.2)$$

with $\Omega_1/\Omega_2 = p/q$. We observe now that we can assume p and q to be relatively prime numbers since $V_{(1)}$ should generate a $U(1)$. In other words Ω_1/Ω_2 is a rational number. One can now proceed with finding the domain structure of the solution.

Another interesting direction to pursue is to consider the various possible domain structures of solutions with black holes attached to Kaluza-Klein bubbles for space-times which are asymptotically Kaluza-Klein space $\mathbb{R}^{1,D-1-q} \times T^q$. As explored via the rod structure in [21] for asymptotically $\mathbb{R}^{1,4} \times S^1$ and $\mathbb{R}^{1,5} \times S^1$ space-times solving vacuum Einstein equations, this could lead to interesting new possibilities for event horizon topologies.

We remark that the vacuum Einstein equations for solutions with $D - 3$ Killing vector fields have an enhanced symmetry, following the construction [22]. The vacuum Einstein equations can be written as a three-dimensional sigma-model with the target space being an $SL(D - 2, \mathbb{R})$ group manifold. This is relevant for asymptotically flat solutions of vacuum Einstein equations in six and seven dimensions. It would be interesting to understand if one can find similar hidden symmetries in the Einstein equations for less number of Killing vectors. This could potentially lead to algebraic solution generating techniques for $D > 5$ similar to the one proposed in [23] for $D = 5$.

Finally, as remarked previously, the domain structure can be generalized to black hole space-times which are not asymptotically flat, including asymptotically Anti-de Sitter space-times. This will be considered in a future publication [9].

Acknowledgments

We thank Pau Figueras for useful discussions. We thank the Carlsberg foundation for support. We thank the Galileo Galilei Institute at Firenze, Italy, the Summer Institute 2008 at Mt. Fuji, Japan, the black hole workshop at Veli Losinj, Croatia, the CERN TH Institute program on black holes, the Banff International Research Station, Canada, and Perugia University at Perugia, Italy, for warm hospitality while this project was carried out.

A Einstein equations for space-times with Killing vector fields

In this appendix we give first a general expression for the Ricci tensor for D -dimensional space-times with p commuting Killing vector fields. We then use this to write down the vacuum Einstein equations for D -dimensional space-times with p commuting Killing vector fields. Finally we examine under which conditions the metric can be written in a block diagonal form with the Killing part of the metric being orthogonal to the rest of the metric.

Ricci tensor for space-times with Killing vector fields

We consider here a given D -dimensional manifold $\mathcal{M}_D$ with a metric with p commuting linearly independent Killing vector fields $V_{(i)}$, $i = 0, 1, \dots, p-1$. Define $n = D - p$. We can always find a coordinate system $x^0, x^1, \dots, x^{p-1}, y^1, \dots, y^n$ such that in this coordinate system the Killing vectors are of the form (2.1) and the metric is of the form (2.2), where G_{ij} , A_a^i and $\tilde{g}_{ab}$ only depend on y^a . Define $K^2 = |\det G_{ij}|$, $\tilde{g} = |\det \tilde{g}_{ab}|$ and $F_{ab}^i = \partial_a A_b^i - \partial_b A_a^i$. The components of the Ricci tensor are

$$R_{ij} = -\frac{1}{2}\partial_a \partial^a G_{ij} - \frac{1}{2}\partial_a (\log K + \log \sqrt{\tilde{g}}) \partial^a G_{ij} + \frac{1}{2}\partial^a G_{ik} G^{kl} \partial_b G_{lj} \\ + \frac{1}{4}G_{ik} G_{jl} \tilde{g}^{ac} \tilde{g}^{bd} F_{ab}^k F_{cd}^l \quad (\text{A.1})$$

$$R_{ia} = R_{ij} A_a^j + \frac{1}{2K\sqrt{\tilde{g}}} \tilde{g}_{ab} \partial_c (K \sqrt{\tilde{g}} G_{ij} \tilde{g}^{bd} \tilde{g}^{ce} F_{de}^j) \quad (\text{A.2})$$

$$R_{ab} = -R_{ij} A_a^i A_b^j + R_{ia} A_b^i + R_{ib} A_a^i - \frac{1}{2} \tilde{g}^{cd} G_{ij} F_{ac}^i F_{bd}^j \\ + \tilde{R}_{ab} - \tilde{D}_a \tilde{D}_b \log K - \frac{1}{4} \text{Tr}(G^{-1} \partial_a G G^{-1} \partial_b G) \quad (\text{A.3})$$

with $\tilde{D}_a \tilde{D}_b \log K = \partial_a \partial_b \log K - \tilde{\Gamma}_{ab}^c \partial_c \log K$ where $\tilde{\Gamma}_{ab}^c$ is the Christoffel symbol as computed from the $\tilde{g}_{ab}$ metric. Define $(C_a)^i_j = G^{ik} \partial_a G_{kj}$. We can write the vacuum Einstein equations $R_{\mu\nu} = 0$ as

$$\partial_a (K \sqrt{\tilde{g}} \tilde{g}^{ab} (C_b)^i_j) = \frac{1}{2} K \sqrt{\tilde{g}} F_{ab}^i G_{jk} \tilde{g}^{ac} \tilde{g}^{bd} F_{cd}^k \quad (\text{A.4})$$

$$\partial_a (K \sqrt{\tilde{g}} G_{ij} \tilde{g}^{ac} \tilde{g}^{bd} F_{cd}^j) = 0 \quad (\text{A.5})$$

$$\tilde{R}_{ab} = \frac{1}{4} \text{Tr}(C_a C_b) + \tilde{D}_a \tilde{D}_b \log K + \frac{1}{2} \tilde{g}^{cd} G_{ij} F_{ac}^i F_{bd}^j \quad (\text{A.6})$$

Results on mixed part of metric solving vacuum Einstein equations

We examine in this section under which conditions one can turn off the A_a^i fields in the general expression for a metric (2.2) solving the vacuum Einstein equations (A.4)-(A.6). The A_a^i field corresponds to the mixed part of the metric (2.2) having indices both in the x^i and y^a directions.

Theorem A.1 *Consider a solution of the vacuum Einstein equations with p commuting Killing vector fields $V_{(i)}$, $i = 0, 1, \dots, p-1$. If the tensors $V_{(0)}^{[\mu_1} V_{(1)}^{\mu_2} \dots V_{(p-1)}^{\mu_p} D^\nu V_{(i)}^{\rho]} = 0$ for all $i = 0, 1, \dots, p-1$ then we can find coordinates such that the metric is of the form*

$$ds^2 = G_{ij} dx^i dx^j + \tilde{g}_{ab} dy^a dy^b \quad (\text{A.7})$$

with the Killing vector fields given by Eq. (2.1).

Proof: Define the one-forms $\xi_\mu^{(i)} = g_{\mu\nu} V_{(i)}^\nu$ for $i = 0, 1, \dots, p-1$. These one-forms span a p -dimensional linear space T^* . Since $V_{(0)}^{[\mu_1} V_{(1)}^{\mu_2} \dots V_{(p-1)}^{\mu_p} D^\nu V_{(i)}^{\rho]} = 0$ we see that $\xi^{(0)} \wedge \xi^{(1)} \wedge \dots \wedge d\xi^{(i)} = 0$ for all $i = 0, 1, \dots, p-1$. This means that for any $\xi \in T^*$ we can find one-forms $\psi^{(i)}$, $i = 0, 1, \dots, p-1$, such that $d\xi = \sum_{i=0}^{p-1} \psi^{(i)} \wedge \xi^{(i)}$. Consider now the $n = D - p$

dimensional tangent space at each point defined by being orthogonal to all one-forms in T^* . From Frobenius' theorem we get that this collection of tangent spaces admits integrable n -dimensional submanifolds. Hence we can find coordinates such that the metric is of the form (A.7). $\square$

To get another perspective on Theorem A.1 we introduce for a given solution of the vacuum Einstein equations with p commuting Killing vector fields $V_{(i)}$ the $(n-2)$ -forms B_i , $i = 0, 1, \dots, p-1$, as

$$(B_i)_{\mu_1 \dots \mu_{n-2}} = \sqrt{g} \epsilon_{\mu_1 \dots \mu_{n-2} \nu_1 \dots \nu_p \rho \sigma} V_{(0)}^{\nu_1} V_{(1)}^{\nu_2} \dots V_{(p-1)}^{\nu_p} D^\rho V_{(i)}^\sigma \quad (\text{A.8})$$

where the ϵ is the D -dimensional ϵ symbol and g is the numerical value of the determinant of the D -dimensional metric. In the (x^i, y^a) coordinates of Eq. (2.2) we see that the μ_j indices only can take values in the y^a directions. We compute now $(B_i)_{a_1 \dots a_{n-2}} = \frac{1}{2} K \sqrt{g} \epsilon_{a_1 \dots a_{n-2} bc} \tilde{g}^{bd} \tilde{g}^{ce} G_{ij} F_{de}^j$ where the ϵ is the n -dimensional ϵ symbol and where $a_j, b, c = 1, \dots, n$. From this we see that $V_{(0)}^{\mu_1} V_{(1)}^{\mu_2} \dots V_{(p-1)}^{\mu_p} D^\nu V_{(i)}^\rho = 0$ if and only if $F_{ab}^i = 0$. Therefore Theorem A.1 tells us that if $F_{ab}^i = 0$ then we can find a gauge transformation such that $A_a^i = 0$. This is already clear locally but Frobenius' theorem ensures that it is also true globally. Another important property of the $(n-2)$ -forms B_i is the following, which follows from the above and Eq. (A.5).

Lemma A.2 *Consider a solution of the vacuum Einstein equations with p commuting Killing vector fields $V_{(i)}$, $i = 0, 1, \dots, p-1$. Then the $(n-2)$ -forms defined in (A.8) are closed $dB_i = 0$. $\square$*

Using this lemma we can prove the following theorem

Theorem A.3 *Consider a solution of the vacuum Einstein equations with $D-2$ commuting Killing vector fields $V_{(i)}$, $i = 0, 1, \dots, D-3$. If the tensor $V_{(0)}^{\mu_1} V_{(1)}^{\mu_2} \dots V_{(D-3)}^{\mu_{D-2}} D^\nu V_{(i)}^\rho = 0$ vanishes at at least one point of the manifold for any given $i = 0, 1, 2, \dots, D-3$ then we can write the metric of the solution in the form (A.7) with the Killing vector fields given by (2.1).*

Proof: This follows from Theorem A.1 and Lemma A.2 since in this case B_i are scalar fields and hence it follows from $dB_i = 0$ for any given $i = 1, 2, \dots, D-2$ that B_i is constant on the manifold. $\square$

This theorem is due to Wald in his book [24] and has been generalized to any dimension by Emparan and Reall in [4]. Thus for $p = D-2$ it is enough that $V_{(0)}^{\mu_1} V_{(1)}^{\mu_2} \dots V_{(D-3)}^{\mu_{D-2}} D^\nu V_{(i)}^\rho$ vanishes at single points for getting the form (A.7) of the metric. Instead for $p < D-2$ we cannot write a generic solution of the vacuum Einstein equations with p commuting Killing vector fields in the form (A.7).

B Parameterizations of topologies from domain structure

We consider the topologies that one can infer from a number of circles fibred over a domain such that the circles shrinks to zero at different points on the boundary of the domain.

Four-sphere topology: We consider here two circles parameterized by ϕ_1 and ϕ_2 fibred over a domain with the shape of the area between the two branches of a parabola taken here to be $z^2 = (z^1)^2$ and furthermore $z^2 \leq 1$. Write the embedding of a four-sphere as

$$x^1 + ix^2 = \sin \theta e^{i\phi_1}, \quad x^3 + ix^4 = \cos \theta \sin \psi e^{i\phi_2}, \quad x^5 = \cos \theta \cos \psi \quad (\text{B.1})$$

where $0 \leq \theta \leq \pi/2$ and $0 \leq \psi \leq \pi$. We then parameterize the domain as

$$z^1 = \cos \theta \cos \psi, \quad z^2 = \cos^2 \theta \quad (\text{B.2})$$

We see that the ϕ_1 circle shrinks to zero for the part of the boundary where $z^2 = 1$ while the ϕ_2 circle shrinks to zero for the part where $z^2 = (z^1)^2$.

Three-sphere topology: We consider here a circle parameterized by ϕ_1 fibred over a domain with the shape of a disc $(z^1)^2 + (z^2)^2 \leq 1$. Write the embedding of a three-sphere as

$$x^1 + ix^2 = \cos \theta e^{i\phi_1}, \quad x^3 + ix^4 = \sin \theta e^{i\phi} \quad (\text{B.3})$$

where $0 \leq \theta \leq \pi/2$. We then parameterize the domain as

$$z^1 = \sin \theta \cos \phi, \quad z^2 = \sin \theta \sin \phi \quad (\text{B.4})$$

We see that the ϕ_1 circle shrinks to zero at the boundary of the disc corresponding to $\theta = \pi/2$ while in the center of the disc the ϕ circle shrinks to zero.

Two-sphere topology: We consider here a domain with the shape of a disc $(z^1 - z_0^1)^2 + (z^2 - z_0^2)^2 \leq 1$. Write the embedding of the two-sphere as

$$x^1 + ix^2 = \cos \theta e^{i\phi}, \quad x^3 = \sin \theta \quad (\text{B.5})$$

where $0 \leq \theta \leq \pi$. We then parameterize the domain as

$$z^1 = z_0^1 + \cos \theta \cos \phi, \quad z^2 = z_0^2 + \cos \theta \sin \phi \quad (\text{B.6})$$

This domain has two sheets: One sheet corresponding to $0 \leq \theta \leq \pi/2$ (i.e. when $x^3 \geq 0$) and the other corresponding to $\pi/2 < \theta \leq \pi$ (i.e. when $x^3 < 0$).

Five-sphere topology: We consider here three circles parameterized by ϕ_1 , ϕ_2 and ϕ_3 fibred over a filled triangle $\frac{3}{2}|z^1| - 1 \leq z^2 \leq \frac{1}{2}$. Write the embedding of the five-sphere as

$$x^1 + ix^2 = \sin \theta e^{i\phi_1}, \quad x^3 + ix^4 = \cos \theta \sin \psi e^{i\phi_2}, \quad x^5 + ix^6 = \cos \theta \cos \psi e^{i\phi_3} \quad (\text{B.7})$$

where $0 \leq \theta, \psi \leq \pi/2$. We then parameterize the domain as

$$z^1 = \cos^2 \theta \cos 2\psi, \quad z^2 = \frac{3}{2} \cos^2 \theta - 1 \quad (\text{B.8})$$

We see that the ϕ_1 circle shrinks to zero at the side of the triangle with $z^2 = \frac{1}{2}$. Instead the ϕ_2 circle shrinks to zero at the side with $z^2 = \frac{3}{2}z^1 - 1$ while the ϕ_3 circle shrinks to zero at the side with $z^2 = -\frac{3}{2}z^1 - 1$.

References

- [1] W. Israel, *Phys. Rev.* **164** (1967) 1776–1779. B. Carter, *Phys. Rev. Lett.* **26** (1971) 331–333. S. W. Hawking, *Commun. Math. Phys.* **25** (1972) 152–166. D. C. Robinson, *Phys. Rev. Lett.* **34** (1975) 905–906.
- [2] R. Emparan and H. S. Reall, *Phys. Rev. Lett.* **88** (2002) 101101, [arXiv:hep-th/0110260](#).
- [3] T. Harmark, *Phys. Rev.* **D70** (2004) 124002, [arXiv:hep-th/0408141](#). T. Harmark and P. Olesen, *Phys. Rev.* **D72** (2005) 124017, [arXiv:hep-th/0508208](#).
- [4] R. Emparan and H. S. Reall, *Phys. Rev.* **D65** (2002) 084025, [arXiv:hep-th/0110258](#).
- [5] S. Hollands and S. Yazadjiev, *Commun. Math. Phys.* **283** (2008) 749–768, [arXiv:0707.2775 \[gr-qc\]](#).
- [6] S. Hollands and S. Yazadjiev, *Class. Quant. Grav.* **25** (2008) 095010, [arXiv:0711.1722 \[gr-qc\]](#). S. Hollands and S. Yazadjiev, [arXiv:0812.3036 \[gr-qc\]](#).
- [7] G. J. Galloway and R. Schoen, *Commun. Math. Phys.* **266** (2006) 571–576, [arXiv:gr-qc/0509107](#).
- [8] S. Hollands, A. Ishibashi, and R. M. Wald, *Commun. Math. Phys.* **271** (2007) 699–722, [arXiv:gr-qc/0605106](#). V. Moncrief and J. Isenberg, *Class. Quant. Grav.* **25** (2008) 195015, [arXiv:0805.1451 \[gr-qc\]](#).
- [9] Work in progress.
- [10] R. C. Myers and M. J. Perry, *Ann. Phys.* **172** (1986) 304.
- [11] R. Emparan, T. Harmark, V. Niarchos, N. A. Obers, and M. J. Rodriguez, *JHEP* **10** (2007) 110, [arXiv:0708.2181 \[hep-th\]](#).
- [12] R. Emparan, T. Harmark, V. Niarchos, and N. A. Obers, *Phys. Rev. Lett.* **102** (2009) 191301, [arXiv:0902.0427 \[hep-th\]](#).
- [13] B. Kleihaus, J. Kunz, and E. Radu, [arXiv:0904.2723 \[hep-th\]](#).
- [14] H. Elvang and P. Figueras, *JHEP* **05** (2007) 050, [arXiv:hep-th/0701035](#).
- [15] H. Iguchi and T. Mishima, *Phys. Rev.* **D75** (2007) 064018, [arXiv:hep-th/0701043](#). J. Evslin and C. Krishnan, *Class. Quant. Grav.* **26** (2009) 125018, [arXiv:0706.1231 \[hep-th\]](#).
- [16] K. Izumi, *Prog. Theor. Phys.* **119** (2008) 757–774, [arXiv:0712.0902 \[hep-th\]](#). H. Elvang and M. J. Rodriguez, *JHEP* **04** (2008) 045, [arXiv:0712.2425 \[hep-th\]](#).
- [17] J. Evslin, *JHEP* **09** (2008) 004, [arXiv:0806.3389 \[hep-th\]](#). Y. Chen and E. Teo, *Phys. Rev.* **D78** (2008) 064062, [arXiv:0808.0587 \[gr-qc\]](#).
- [18] Y. Morisawa and D. Ida, *Phys. Rev.* **D69** (2004) 124005, [arXiv:gr-qc/0401100](#).
- [19] R. Emparan, *JHEP* **03** (2004) 064, [arXiv:hep-th/0402149](#).
- [20] K. Copsey and G. T. Horowitz, *Phys. Rev.* **D73** (2006) 024015, [arXiv:hep-th/0505278](#).
- [21] H. Elvang, T. Harmark, and N. A. Obers, *JHEP* **01** (2005) 003, [hep-th/0407050](#).
- [22] D. Maison, *Gen. Rel. Grav.* **10** (1979) 717–723.
- [23] S. Giusto and A. Saxena, *Class. Quant. Grav.* **24** (2007) 4269–4294, [arXiv:0705.4484 \[hep-th\]](#).
- [24] R. M. Wald, *General Relativity*. The University of Chicago Press, 1984.